

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
22 March 2001 (22.03.2001)

PCT

(10) International Publication Number
WO 01/19828 A2

(51) International Patent Classification⁷: **C07D 487/04**,
471/04; 491/04, 498/04, A61K 31/519, 31/5025, 31/4353,
31/551, A61P 35/00 // (C07D 487/04, 239:00, 209:00)
(C07D 487/04, 237:00, 209:00) (C07D 471/04, 231:00,
221:00) (C07D 487/04, 243:00, 239:00)

(21) International Application Number: PCT/US00/25357

(22) International Filing Date:
15 September 2000 (15.09.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/154,618 17 September 1999 (17.09.1999) US

(74) Agents: **WAGNER, Richard, W.** et al.; Hamilton, Brook,
Smith & Reynolds, P.C., Two Militia Drive, Lexington, MA
02421 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,
TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG,
CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*): **BASF**
AKTIENGESELLSCHAFT [DE/DE]; 67056 Lud-
wigshafen (DE).

(72) Inventor; and

(75) Inventor/Applicant (*for US only*): **HIRST, Gavin, C.**
[GB/US]; 112 Robert Road, Marlboro, MA 01752 (US).

Published:

— *Without international search report and to be republished
upon receipt of that report.*

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

WO 01/19828 A2

(54) Title: KINASE INHIBITORS AS THERAPEUTIC AGENTS

(57) Abstract: The present invention is directed to a compound of Formula (I) as defined herein which are useful as kinase inhibitors.

KINASE INHIBITORS AS THERAPEUTIC AGENTS

RELATED APPLICATION

This application claims the benefit of United States Provisional Application
5 No. 60/154,618, filed September 17, 1999, the entire teachings of which are
incorporated herein by reference.

BACKGROUND OF THE INVENTION

There are at least 400 enzymes identified as protein kinases. These enzymes
10 catalyze the phosphorylation of target protein substrates. The phosphorylation is
usually a transfer reaction of a phosphate group from ATP to the protein substrate.
The specific structure in the target substrate to which the phosphate is transferred is
a tyrosine, serine or threonine residue. Since these amino acid residues are the target
structures for the phosphoryl transfer, these protein kinase enzymes are commonly
15 referred to as tyrosine kinases or serine/threonine kinases.

The phosphorylation reactions, and counteracting phosphatase reactions, at
the tyrosine, serine and threonine residues are involved in countless cellular
processes that underlie responses to diverse intracellular signals (typically mediated
through cellular receptors), regulation of cellular functions, and activation or
20 deactivation of cellular processes. A cascade of protein kinases often participate in
intracellular signal transduction and are necessary for the realization of these cellular
processes. Because of their ubiquity in these processes, the protein kinases can be
found as an integral part of the plasma membrane or as cytoplasmic enzymes or
localized in the nucleus, often as components of enzyme complexes. In many
25 instances, these protein kinases are an essential element of enzyme and structural
protein complexes that determine where and when a cellular process occurs within a
cell.

Protein Tyrosine Kinases. Protein tyrosine kinases (PTKs) are enzymes
which catalyze the phosphorylation of specific tyrosine residues in cellular proteins.
30 This post-translational modification of these substrate proteins, often enzymes
themselves, acts as a molecular switch regulating cell proliferation, activation or

-2-

differentiation (for review, see Schlessinger and Ulrich, 1992, *Neuron* 9:383-391). Aberrant or excessive PTK activity has been observed in many disease states including benign and malignant proliferative disorders as well as diseases resulting from inappropriate activation of the immune system (e.g., autoimmune disorders),
5 allograft rejection, and graft vs. host disease. In addition, endothelial-cell specific receptor PTKs such as KDR and Tie-2 mediate the angiogenic process, and are thus involved in supporting the progression of cancers and other diseases involving inappropriate vascularization (e.g., diabetic retinopathy, choroidal
10 retinopathy of prematurity, infantile hemangiomas).

Tyrosine kinases can be of the receptor-type (having extracellular, transmembrane and intracellular domains) or the non-receptor type (being wholly intracellular).

Receptor Tyrosine Kinases (RTKs). The RTKs comprise a large family of
15 transmembrane receptors with diverse biological activities. At present, at least nineteen (19) distinct RTK subfamilies have been identified. The receptor tyrosine kinase (RTK) family includes receptors that are crucial for the growth and differentiation of a variety of cell types (Yarden and Ullrich, *Ann. Rev. Biochem.* 57:433-478, 1988; Ullrich and Schlessinger, *Cell* 61:243-254, 1990). The intrinsic
20 function of RTKs is activated upon ligand binding, which results in phosphorylation of the receptor and multiple cellular substrates, and subsequently in a variety of cellular responses (Ullrich & Schlessinger, 1990, *Cell* 61:203-212). Thus, receptor tyrosine kinase mediated signal transduction is initiated by extracellular interaction with a specific growth factor (ligand), typically followed by receptor dimerization,
25 stimulation of the intrinsic protein tyrosine kinase activity and receptor trans-phosphorylation. Binding sites are thereby created for intracellular signal transduction molecules and lead to the formation of complexes with a spectrum of cytoplasmic signaling molecules that facilitate the appropriate cellular response. (e.g., cell division, differentiation, metabolic effects, changes in the extracellular
30 microenvironment) see Schlessinger and Ullrich, 1992, *Neuron* 9:1-20.

Proteins with SH2 (src homology -2) or phosphotyrosine binding (PTB) domains bind activated tyrosine kinase receptors and their substrates with high affinity to propagate signals into cell. Both of the domains recognize phosphotyrosine. (Fantl *et al.*, 1992, *Cell* 69:413-423; Songyang *et al.*, 1994, *Mol. Cell. Biol.* 14:2777-2785; Songyang *et al.*, 1993, *Cell* 72:767-778; and Koch *et al.*, 1991, *Science* 252:668-678; Shoelson, *Curr. Opin. Chem. Biol.* (1997), 1(2), 227-234; Cowburn, *Curr. Opin. Struct. Biol.* (1997), 7(6), 835-838). Several intracellular substrate proteins that associate with receptor tyrosine kinases (RTKs) have been identified. They may be divided into two principal groups: (1) substrates which have a catalytic domain; and (2) substrates which lack such a domain but serve as adapters and associate with catalytically active molecules (Songyang *et al.*, 1993, *Cell* 72:767-778). The specificity of the interactions between receptors or proteins and SH2 or PTB domains of their substrates is determined by the amino acid residues immediately surrounding the phosphorylated tyrosine residue. For example, differences in the binding affinities between SH2 domains and the amino acid sequences surrounding the phosphotyrosine residues on particular receptors correlate with the observed differences in their substrate phosphorylation profiles (Songyang *et al.*, 1993, *Cell* 72:767-778). Observations suggest that the function of each receptor tyrosine kinase is determined not only by its pattern of expression and ligand availability but also by the array of downstream signal transduction pathways that are activated by a particular receptor as well as the timing and duration of those stimuli. Thus, phosphorylation provides an important regulatory step which determines the selectivity of signaling pathways recruited by specific growth factor receptors, as well as differentiation factor receptors.

Several receptor tyrosine kinases such as FGFR-1, PDGFR, TIE-2 and c-Met, and growth factors that bind thereto, have been suggested to play a role in angiogenesis, although some may promote angiogenesis indirectly (Mustonen and Alitalo, *J. Cell Biol.* 129:895-898, 1995). One such receptor tyrosine kinase, known as "fetal liver kinase 1" (FLK-1), is a member of the type III subclass of RTKs. An alternative designation for human FLK-1 is "kinase insert domain-containing

-4-

receptor" (KDR) (Terman *et al.*, *Oncogene* 6:1677-83, 1991). Another alternative designation for FLK-1/KDR is "vascular endothelial cell growth factor receptor 2" (VEGFR-2) since it binds VEGF with high affinity. The murine version of FLK-1/VEGFR-2 has also been called NYK (Oelrichs *et al.*, *Oncogene* 8(1):11-15, 1993).

5 DNAs encoding mouse, rat and human FLK-1 have been isolated, and the nucleotide and encoded amino acid sequences reported (Matthews *et al.*, *Proc. Natl. Acad. Sci. USA*, 88:9026-30, 1991; Terman *et al.*, 1991, *supra*; Terman *et al.*, *Biochem. Biophys. Res. Comm.* 187:1579-86, 1992; Sarzani *et al.*, *supra*; and Millauer *et al.*, *Cell* 72:835-846, 1993). Numerous studies such as those reported in
10 Millauer *et al.*, *supra*, suggest that VEGF and FLK-1/KDR/VEGFR-2 are a ligand-receptor pair that play an important role in the proliferation of vascular endothelial cells, and formation and sprouting of blood vessels, termed vasculogenesis and angiogenesis, respectively.

Another type III subclass RTK designated "fms-like tyrosine kinase-1" (Flt-1) is related to FLK-1/KDR (DeVries *et al.* *Science* 255:989-991, 1992; Shibuya *et al.*, *Oncogene* 5:519-524, 1990). An alternative designation for Flt-1 is "vascular endothelial cell growth factor receptor 1" (VEGFR-1). To date, members of the FLK-1/ KDR/VEGFR-2 and Flt-1/ VEGFR-1 subfamilies have been found expressed primarily on endothelial cells. These subclass members are specifically
20 stimulated by members of the vascular endothelial cell growth factor (VEGF) family of ligands (Klagsburn and D'Amore, *Cytokine & Growth Factor Reviews* 7: 259-270, 1996). Vascular endothelial cell growth factor (VEGF) binds to Flt-1 with higher affinity than to FLK-1/KDR and is mitogenic toward vascular endothelial cells (Terman *et al.*, 1992, *supra*; Mustonen *et al.* *supra*; DeVries *et al.*, *supra*). Flt-1
25 is believed to be essential for endothelial organization during vascular development. Flt-1 expression is associated with early vascular development in mouse embryos, and with neovascularization during wound healing (Mustonen and Alitalo, *supra*). Expression of Flt-1 in monocytes, osteoclasts, and osteoblasts, as well as in adult tissues such as kidney glomeruli suggests an additional function for
30 this receptor that is not related to cell growth (Mustonen and Alitalo, *supra*).

-5-

As previously stated, recent evidence suggests that VEGF plays a role in the stimulation of both normal and pathological angiogenesis (Jakeman *et al.*, *Endocrinology* 133: 848-859, 1993; Kolch *et al.*, *Breast Cancer Research and Treatment* 36: 139-155, 1995; Ferrara *et al.*, *Endocrine Reviews* 18(1); 4-25, 1997; Ferrara *et al.*, Regulation of Angiogenesis (ed. L. D. Goldberg and E.M. Rosen), 209-232, 1997). In addition, VEGF has been implicated in the control and enhancement of vascular permeability (Connolly, *et al.*, *J. Biol. Chem.* 264: 20017-20024, 1989; Brown *et al.*, Regulation of Angiogenesis (ed. L.D. Goldberg and E.M. Rosen), 233-269, 1997). Different forms of VEGF arising from alternative splicing of mRNA have been reported, including the four species described by Ferrara *et al.* (*J. Cell. Biochem.* 47:211-218, 1991). Both secreted and predominantly cell-associated species of VEGF have been identified by Ferrara *et al. supra*, and the protein is known to exist in the form of disulfide linked dimers.

Several related homologs of VEGF have recently been identified. However, their roles in normal physiological and disease processes have not yet been elucidated. In addition, the members of the VEGF family are often coexpressed with VEGF in a number of tissues and are, in general, capable of forming heterodimers with VEGF. This property likely alters the receptor specificity and biological effects of the heterodimers and further complicates the elucidation of their specific functions as illustrated below (Korpelainen and Alitalo, *Curr. Opin. Cell Biol.*, 159-164, 1998 and references cited therein).

Placenta growth factor (PlGF) has an amino acid sequence that exhibits significant homology to the VEGF sequence (Park *et al.*, *J. Biol. Chem.* 269:25646-54, 1994; Maglione *et al. Oncogene* 8:925-31, 1993). As with VEGF, different species of PlGF arise from alternative splicing of mRNA, and the protein exists in dimeric form (Park *et al.*, *supra*). PlGF-1 and PlGF-2 bind to Flt-1 with high affinity, and PlGF-2 also avidly binds to neuropilin-1 (Migdal *et al.*, *J. Biol. Chem.* 273 (35): 22272-22278), but neither binds to FLK-1/KDR (Park *et al.*, *supra*). PlGF has been reported to potentiate both the vascular permeability and mitogenic effect of VEGF on endothelial cells when VEGF is present at low concentrations (purportedly due to heterodimer formation) (Park *et al.*, *supra*).

VEGF-B is produced as two isoforms (167 and 185 residues) that also appear to bind Flt-1/VEGFR-1. It may play a role in the regulation of extracellular matrix degradation, cell adhesion, and migration through modulation of the expression and activity of urokinase type plasminogen activator and plasminogen activator inhibitor
5 1 (Pepper *et al*, *Proc. Natl. Acad. Sci. U. S. A.* (1998), 95(20): 11709-11714).

VEGF-C was originally cloned as a ligand for VEGFR-3/Flt-4 which is primarily expressed by lymphatic endothelial cells. In its fully processed form, VEGF-C can also bind KDR/VEGFR-2 and stimulate proliferation and migration of endothelial cells *in vitro* and angiogenesis in *in vivo* models (Lymboussaki *et al*,
10 *Am. J. Pathol.* (1998), 153(2): 395-403; Witzienbichler *et al*, *Am. J. Pathol.* (1998), 153(2), 381-394). The transgenic overexpression of VEGF-C causes proliferation and enlargement of only lymphatic vessels, while blood vessels are unaffected. Unlike VEGF, the expression of VEGF-C is not induced by hypoxia (Ristimaki *et al*, *J. Biol. Chem.* (1998), 273(14),8413-8418).

15 The most recently discovered VEGF-D is structurally very similar to VEGF-C. VEGF-D is reported to bind and activate at least two VEGFRs, VEGFR-3/Flt-4 and KDR/VEGFR-2. It was originally cloned as a c-fos inducible mitogen for fibroblasts and is most prominently expressed in the mesenchymal cells of the lung and skin (Achen *et al*, *Proc. Natl. Acad. Sci. U. S. A.* (1998), 95(2), 548-553 and
20 references therein).

As for VEGF, VEGF-C and VEGF-D have been claimed to induce increases in vascular permeability *in vivo* in a Miles assay when injected into cutaneous tissue (PCT/US97/14696; WO98/07832, Witzienbichler *et al.*, *supra*). The physiological role and significance of these ligands in modulating vascular hyperpermeability and
25 endothelial responses in tissues where they are expressed remains uncertain.

There has been recently reported a virally encoded, novel type of vascular endothelial growth factor, VEGF-E (NZ-7 VEGF), which preferentially utilizes KDR/Flk-1 receptor and carries a potent mitotic activity without heparin-binding domain (Meyer *et al*, *EMBO J.* (1999), 18(2), 363-374; Ogawa *et al*, *J. Biol. Chem.*
30 (1998), 273(47), 31273-31282.). VEGF-E sequences possess 25% homology to mammalian VEGF and are encoded by the parapoxvirus Orf virus (OV). This

-7-

parapoxvirus that affects sheep and goats and occasionally, humans, to generate lesions with angiogenesis. VEGF-E is a dimer of about 20 kDa with no basic domain nor affinity for heparin, but has the characteristic cysteine knot motif present in all mammalian VEGFs, and was surprisingly found to possess potency and bioactivities similar to the heparin-binding VEGF165 isoform of VEGF-A, i.e. both factors stimulate the release of tissue factor (TF), the proliferation, chemotaxis and sprouting of cultured vascular endothelial cells in vitro and angiogenesis in vivo. Like VEGF165, VEGF-E was found to bind with high affinity to VEGF receptor-2 (KDR) resulting in receptor autophosphorylation and a biphasic rise in free intracellular Ca²⁺ concentrations, while in contrast to VEGF165, VEGF-E did not bind to VEGF receptor-1 (Flt-1).

Based upon emerging discoveries of other homologs of VEGF and VEGFRs and the precedents for ligand and receptor heterodimerization, the actions of such VEGF homologs may involve formation of VEGF ligand heterodimers, and/or heterodimerization of receptors, or binding to a yet undiscovered VEGFR (Witzenbichler *et al.*, *supra*). Also, recent reports suggest neuropilin-1 (Migdal *et al.*, *supra*) or VEGFR-3/Flt-4 (Witzenbichler *et al.*, *supra*), or receptors other than KDR/VEGFR-2 may be involved in the induction of vascular permeability (Stacker, S.A., Vitali, A., Domagala, T., Nice, E., and Wilks, A.F., "Angiogenesis and Cancer" Conference, Amer. Assoc. Cancer Res., Jan. 1998, Orlando, FL; Williams, *Diabetologia* 40: S118-120 (1997)). Until now, no direct evidence for the essential role of KDR in VEGF-mediated vascular hyperpermeability has been disclosed.

The Non-Receptor Tyrosine Kinases. The non-receptor tyrosine kinases represent a collection of cellular enzymes which lack extracellular and transmembrane sequences. At present, over twenty-four individual non-receptor tyrosine kinases, comprising eleven (11) subfamilies (Src, Frk, Btk, Csk, Abl, Zap70, Fes/Fps, Fak, Jak, Ack and LIMK) have been identified. At present, the Src subfamily of non-receptor tyrosine kinases is comprised of the largest number of PTKs and include Src, Yes, Fyn, Lyn, Lck, Blk, Hck, Fgr and Yrk. The Src

30

subfamily of enzymes has been linked to oncogenesis and immune responses. A more detailed discussion of non-receptor tyrosine kinases is provided in Bolen, 1993, *Oncogene* 8:2025-2031, which is incorporated herein by reference.

Many of the tyrosine kinases, whether an RTK or non-receptor tyrosine
5 kinase, have been found to be involved in cellular signaling pathways involved in numerous pathogenic conditions, including cancer, psoriasis, and other hyperproliferative disorders or hyper-immune responses.

Development of Compounds to Modulate the PTKs. In view of the surmised importance of PTKs to the control, regulation, and modulation of cell proliferation,
10 the diseases and disorders associated with abnormal cell proliferation, many attempts have been made to identify receptor and non-receptor tyrosine kinase "inhibitors" using a variety of approaches, including the use of mutant ligands (U.S. Application No. 4,966,849), soluble receptors and antibodies (Application No. WO 94/10202; Kendall & Thomas, 1994, *Proc. Natl. Acad. Sci* 90:10705-09; Kim *et al.*,
15 1993, *Nature* 362:841-844), RNA ligands (Jellinek, *et al.*, *Biochemistry* 33:10450-56; Takano, *et al.*, 1993, *Mol. Bio. Cell* 4:358A; Kinsella, *et al.* 1992, *Exp. Cell Res.* 199:56-62; Wright, *et al.*, 1992, *J. Cellular Phys.* 152:448-57) and tyrosine kinase inhibitors (WO 94/03427; WO 92/21660; WO 91/15495; WO 94/14808; U.S. Patent No. 5,330,992; Mariani, *et al.*, 1994, *Proc. Am. Assoc. Cancer Res.* 35:2268).

20 More recently, attempts have been made to identify small molecules which act as tyrosine kinase inhibitors. For example, bis monocyclic, bicyclic or heterocyclic aryl compounds (PCT WO 92/20642) and vinylene-azaindole derivatives (PCT WO 94/14808) have been described generally as tyrosine kinase inhibitors. Styryl compounds (U.S. Patent No. 5,217,999), styryl-substituted pyridyl
25 compounds (U.S. Patent No. 5,302,606), certain quinazoline derivatives (EP Application No. 0 566 266 A1; *Expert Opin. Ther. Pat.* (1998), 8(4): 475-478), selenoindoles and selenides (PCT WO 94/03427), tricyclic polyhydroxylic compounds (PCT WO 92/21660) and benzylphosphonic acid compounds (PCT WO 91/15495) have been described as compounds for use as tyrosine kinase inhibitors
30 for use in the treatment of cancer. Anilino-cinnolines (PCT WO97/34876) and

quinazoline derivative compounds (PCT WO97/22596; PCT WO97/42187) have been described as inhibitors of angiogenesis and vascular permeability.

In addition, attempts have been made to identify small molecules which act as serine/threonine kinase inhibitors. For example, bis(indolylmaleimide) compounds have been described as inhibiting particular PKC serine/threonine kinase isoforms whose signal transducing function is associated with altered vascular permeability in VEGF-related diseases (PCT WO97/40830; PCT WO97/40831).

Plk-1 Kinase Inhibitors

Plk-1 is a serine/threonine kinase which is an important regulator of cell cycle progression. It plays critical roles in the assembly and the dynamic function of the mitotic spindle apparatus. Plk-1 and related kinases have also been shown to be closely involved in the activation and inactivation of other cell cycle regulators, such as cyclin-dependent kinases. High levels of Plk-1 expression are associated with cell proliferation activities. It is often found in malignant tumors of various origins. Inhibitors of Plk-1 are expected to block cancer cell proliferation by disrupting processes involving mitotic spindles and inappropriately activated cyclin-dependent kinases.

Cdc2/Cyclin B Kinase Inhibitors (Cdc2 is also known as cdk1)

Cdc2/cyclin B is another serine/threonine kinase enzyme which belongs to the cyclin-dependent kinase (cdks) family. These enzymes are involved in the critical transition between various phases of cell cycle progression. It is believed that uncontrolled cell proliferation, which is the hallmark of cancer is dependent upon elevated cdk activities in these cells. The inhibition of elevated cdk activities in cancer cells by cdc2/cyclin B kinase inhibitors could suppress proliferation and may restore the normal control of cell cycle progression.


The identification of effective small compounds which specifically inhibit signal transduction and cellular proliferation by modulating the activity of receptor and non-receptor tyrosine and serine/threonine kinases to regulate and modulate abnormal or inappropriate cell proliferation, differentiation, or metabolism is

-10-

therefore desirable. In particular, the identification of methods and compounds that specifically inhibit the function of a tyrosine kinase which is essential for antiogenic processes or the formation of vascular hyperpermeability leading to edema, ascites, effusions, exudates, and macromolecular extravasation and matrix deposition as well as associated disorders would be beneficial.


SUMMARY OF THE INVENTION

The present invention provides a compound of Formula (I), the racemic-diastereomeric mixtures, optical isomers, pharmaceutically-acceptable salts, prodrugs or biologically active metabolites thereof, selected from the group consisting of sub-formulas 1-109:




15


-11-




13




14




15




16




17




18




19




20




21




22




23




24




25



26






27




28

5



-12-





5






-13-





5



-14-




5




10


-15-




81




82



83




84




85




86




87




88




89




90




91




92




93




94




95




96




97



98



99




100

10

-16-




101




102




103




104




105




106




107




108




109



110




111




112


5




113



114




115



116

and




117


10

-17-

R_1 is of the formula



where Z^{100} is or a group optionally substituted with R_b ,
 selected from the group consisting of cycloalkyl, naphthyl,
 tetrahydronaphthyl, benzothienyl, furanyl, thienyl, benzoxazolyl,



benzothiazolyl, , thiazolyl, benzofuranyl,
 2,3-dihydrobenzofuranyl, indolyl, isoxazolyl, tetrahydropyranyl,
 tetrahydrofuranyl, piperidinyl, pyrazolyl, pyrrolyl, oxazolyl, isothiazolyl,
 oxadiazolyl, thiadiazolyl, indolinyl, indazolyl, benzoisothiazolyl, pyrido-
 oxazolyl, pyrido-thiazolyl, pyrimido-oxazolyl, pyrimido-thiazolyl and
 benzimidazolyl;

Z^{110} is a covalent bond, or an optionally substituted (C_1-C_6) which is
 optionally substituted with one or more substituents selected from the group
 consisting of alkyl, CN, OH, halogen, NO_2 , COOH, substituted or
 unsubstituted amino and substituted or unsubstituted phenyl;

Z^{111} is a covalent bond, an optionally substituted (C_1-C_6) or an optionally
 substituted $-(CH_2)_n$ -cycloalkyl- $(CH_2)_n$ -; where the optionally substituted
 groups are optionally substituted with one or more substituents selected from
 the group consisting of alkyl, CN, OH, halogen, NO_2 , COOH, substituted or
 unsubstituted amino and substituted or unsubstituted phenyl;

R_a and R_b each represent one or more substituents for each occurrence
 independently selected from the group consisting of hydrogen, halogen, -CN,
 - NO_2 , -C(O)OH, -C(O)H, -OH, -C(O)O-alkyl, substituted or unsubstituted
 carboxamido, tetrazolyl, trifluoromethylcarbonylamino,

- trifluoromethylsulfonamido, substituted or unsubstituted alkyl, substituted or unsubstituted alkoxy, substituted or unsubstituted aryl, substituted or unsubstituted alkenyl, substituted or unsubstituted aryloxy, substituted or unsubstituted heteroaryloxy, substituted or unsubstituted arylalkyl, substituted or unsubstituted alkynyl, substituted or unsubstituted amino, substituted or unsubstituted aminoalkyl, substituted or unsubstituted amido groups, substituted or unsubstituted heteroarylthio, substituted or unsubstituted arylthio, $-Z^{105}-C(O)N(R)_2$, $-Z^{105}-N(R)-C(O)-Z^{200}$, $-Z^{105}-N(R)-S(O)_2-Z^{200}$, $-Z^{105}-N(R)-C(O)-N(R)-Z^{200}$, R_e and CH_2OR_e ;
- where R_e for each occurrence is independently hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted aryl, $-CH_2-NR_dR_e$, $-W-(CH_2)_t-NR_dR_e$, $-W-(CH_2)_t-Oalkyl$, $-W-(CH_2)_t-S-alkyl$, or $-W-(CH_2)_t-OH$; Z^{105} for each occurrence is independently a covalent bond or (C_1-C_6) ; Z^{200} for each occurrence is independently a substituted or unsubstituted (C_1-C_6) , substituted or unsubstituted phenyl or substituted or unsubstituted $-(C_1-C_6)$ -phenyl;
- R_d and R_e for each occurrence are independently H, alkyl, alkanoyl or SO_2 -alkyl; or R_d , R_e and the nitrogen atom to which they are attached together form a five- or six-membered heterocyclic ring; t for each occurrence is independently an integer from 2 to 6; W for each occurrence is independently a direct bond or O, S, $S(O)$, $S(O)_2$, or NR_f , wherein R_f for each occurrence is independently H or alkyl; or R_f is a substituted or unsubstituted carbocyclic or heterocyclic ring fused with ring 2;
- A is $-O-$; $-S-$; $-S(O)_p-$; $-N(R)-$; $-N(C(O)OR)-$; $-N(C(O)R)-$; $-N(SO_2R)-$; $-CH_2O-$; $-CH_2S-$; $-CH_2N(R)-$; $-CH(NR)-$; $-CH_2N(C(O)R)-$; $-CH_2N(C(O)OR)-$; $-CH_2N(SO_2R)-$; $-CH(NHR)-$; $-CH(NHC(O)R)-$; $-CH(NHSO_2R)-$; $-CH(NHC(O)OR)-$; $-CH(OC(O)R)-$; $-CH(OC(O)NHR)-$; $-CH=CH-$; $-C(=NOR)-$; $-C(O)-$; $-CH(OR)-$; $-C(O)N(R)-$; $-N(R)C(O)-$; $-N(R)S(O)_p-$; $-OC(O)N(R)-$; $-N(R)-C(O)-(CH_2)_n-N(R)-$;

-19-

$-N(R)C(O)O-$; $-N(R)-(CH_2)_{n+1}-C(O)-$; $-S(O)_pN(R)-$;
 $-O-(CR_2)_{n+1}-C(O)-$; $-O-(CR_2)_{n+1}-O-$;
 $-N(C(O)R)S(O)_p-$; $-N(R)S(O)_pN(R)-$; $-N(R)-C(O)-(CH_2)_n-O-$;
 $-C(O)N(R)C(O)-$; $-S(O)_pN(R)C(O)-$; $-OS(O)_pN(R)-$; $-N(R)S(O)_pO-$;
 $-N(R)S(O)_pC(O)-$; $-SO_pN(C(O)R)-$; $-N(R)SO_pN(R)-$; $-C(O)O-$;
 $-N(R)P(OR_g)O-$; $-N(R)P(OR_g)-$; $-N(R)P(O)(OR_g)O-$;
 $-N(R)P(O)(OR_g)-$;
 $-N(C(O)R)P(OR_g)O-$; $-N(C(O)R)P(OR_g)-$; $-N(C(O)R)P(O)(OR_g)O-$,
 or
 $-N(C(O)R)P(OR_g)-$;

where R for each occurrence is independently H, substituted or unsubstituted alkyl, substituted or unsubstituted arylalkyl or substituted or unsubstituted aryl;

R_g for each occurrence is independently H, substituted or unsubstituted alkyl, substituted or unsubstituted arylalkyl, substituted or unsubstituted cycloalkyl or substituted or unsubstituted aryl;

p is 1 or 2;

or in a phosphorus containing group, the nitrogen atom, the phosphorus atom, R and R_g together form a five- or six-membered heterocyclic ring; or

A is $NRSO_2$ and R, R_a and the nitrogen atom together form a substituted or unsubstituted five or six-membered heterocyclic ring fused to ring 1;

R_2 is $-Z^{101}-Z^{102}$;

Z^{101} is a covalent bond, $-(C_1-C_6)-$, $-(C_1-C_6)-O-$, $-(C_1-C_6)-C(O)-$, $-(C_1-C_6)-C(O)O-$, $-(C_1-C_6)-C(O)-NH-$, $-(C_1-C_6)-C(O)-N((C_1-C_6))-$ or a substituted or unsubstituted phenyl group;

Z^{102} is hydrogen, a substituted or unsubstituted alkyl group, a substituted or unsubstituted cycloalkyl group, a substituted or unsubstituted, saturated or unsaturated heterocyclic group, or a substituted or unsubstituted, saturated or unsaturated heterobicyclic group;

-20-

said substituted heterocyclic or substituted heterobicyclic group having one or more substituents each independently selected from the group consisting of hydroxyl, cyano, substituted or unsubstituted alkoxy, substituted or unsubstituted sulfonamido, substituted or unsubstituted ureido, substituted or unsubstituted carboxamido; substituted or unsubstituted amino, oxo, a saturated, unsaturated or aromatic, substituted or unsubstituted heterocyclic group comprising one or more nitrogen atoms, one or more oxygen atoms or a combination thereof;

wherein said nitrogen atoms are independently optionally substituted by a substituted or unsubstituted alkyl, substituted or unsubstituted aryl or substituted or unsubstituted arylalkyl group; or

R_2 is of the formula B-E, wherein B is a substituted or unsubstituted cycloalkyl, substituted or unsubstituted azacycloalkyl, substituted or unsubstituted amino, substituted or unsubstituted aminoalkylsulfonyl, substituted or unsubstituted alkoxyalkyl, substituted or unsubstituted alkoxy, substituted or unsubstituted aminoalkylcarbonyl, hydroxy, substituted or unsubstituted alkylene, substituted or unsubstituted aminoalkyl, substituted or unsubstituted alkylencarbonyl or substituted or unsubstituted aminoalkylcarbonyl group; and E is substituted or unsubstituted azacycloalkyl, substituted or unsubstituted azacycloalkylcarbonyl, substituted or unsubstituted azacycloalkylsulfonyl, substituted or unsubstituted azacycloalkylalkyl, substituted or unsubstituted heteroaryl, substituted or unsubstituted heteroarylcarbonyl, substituted or unsubstituted heteroarylsulfonyl, substituted or unsubstituted heteroarylalkyl, substituted or unsubstituted azacycloalkylcarbonylamino, substituted or unsubstituted heteroarylcarbonylamino or substituted or unsubstituted aryl;

R_3 for each occurrence is independently hydrogen, hydroxy, substituted or unsubstituted alkyl or substituted or unsubstituted alkoxy;

a is 1 and D_1 , G_1 , J_1 , L_1 and M_1 are each independently selected from the group

consisting of CR_a and N, provided that at least two of D_1 , G_1 , J_1 , L_1 and M_1 are CR_a ; or

-21-

a is 0, and one of D_1 , G_1 , L_1 and M_1 is NR_a , one of D_1 , G_1 , L_1 and M_1 is CR_a and the remainder are independently selected from the group consisting of CR_a and N , wherein R_a is as defined above;

b is 1 and D_2 , G_2 , J_2 , L_2 and M_2 are each independently selected from the group

consisting of CR_a and N , provided that at least two of D_2 , G_2 , J_2 , L_2 and M_2 are CR_a ; or

b is 0, and one of D_2 , G_2 , L_2 and M_2 is NR_a , one of D_2 , G_2 , L_2 and M_2 is CR_a and the remainder are independently selected from the group consisting of CR_a and N , wherein R_a is as defined above; and

n for each occurrence is independently an integer from 0 to 6.

Each of the sub-formulas 1-117 is a preferred embodiment of the present application.

The compounds of this invention are useful as inhibitors of serine/threonine and tyrosine kinases. In particular, compounds of this invention are useful as inhibitors of tyrosine kinases that are important in hyperproliferative diseases, especially in cancer and in the process of angiogenesis. For example, certain of these compounds are inhibitors of such receptor kinases as KDR, Flt-1, FGFR, PDGFR, c-Met, TIE-2 or IGF-1-R. Since certain of these compounds are anti-angiogenic, they are important substances for inhibiting the progression of disease states where angiogenesis is an important component. Certain compounds of the invention are effective as inhibitors of such serine/threonine kinases as PKCs, erk, MAP kinases, cdks, Plk-1 or Raf-1. These compounds are useful in the treatment of cancer, and hyperproliferative disorders. In addition, certain compounds are effective inhibitors of non-receptor kinases such as those of the Src (for example, Ick, blk and lyn), Tec, Csk, Jak, Map, Nik and Syk families. These compounds are useful in the treatment of cancer, hyperproliferative disorders and immunologic diseases.

Certain compounds of this invention are selective TIE-2 kinase inhibitors which may be anti-angiogenic (especially in combination with one or more VEGFR inhibitors), or pro-angiogenic, when employed in the presence of, or in conjunction

-22-

with, a VEGF-related stimulus. In this manner such inhibitors can be used in the promotion of therapeutic angiogenesis to treat, for example, ischemia, infarct or occlusion, or to promote wound healing.

5 The present invention provides a method of inhibiting the kinase activity of tyrosine kinases and serine/threonine kinases comprising the administration of a compound represented by formulas 1-109 to said kinase in sufficient concentration to inhibit the enzyme activity of said kinase.

10 The present invention further includes the use of these compounds in pharmaceutical compositions with a pharmaceutically effective amount of the above-described compounds and a pharmaceutically acceptable carrier or excipient. These pharmaceutical compositions can be administered to individuals to slow or halt the process of angiogenesis in angiogenesis-aided diseases, or to treat edema, effusions, exudates or ascites and other conditions associated with vascular hyperpermeability. Certain pharmaceutical compositions can be administered to individuals to treat
15 cancer and hyperproliferative disorders by inhibiting serine/threonine kinases such as cdk, Plk-1, erk, etc.

DETAILED DESCRIPTION OF THE INVENTION

20 Progression through the eukaryotic cell cycle is controlled by a family of kinases called cyclin dependent kinases (CDKs) (Myerson *et al.*, *EMBO Journal*, 11:2909-2917 (1992)). The regulation of CDK activation is complex, but requires the association of the CDK with a member of the cyclin family of regulatory subunits (Draetta, *Trends in Cell Biology*, 3:287-289 (1993)); Murray and Kirschner, *Nature*, 339:275-280 (1989); Solomon *et al.*, *Molecular Biology of the Cell*, 3:13-27
25 (1992)). A further level of regulation occurs through both activating and inactivating phosphorylations of the CDK subunit (Draetta, *Trends in Cell Biology*, 3:287-289 (1993)); Murray and Kirschner, *Nature*, 339:275-280 (1989); Solomon *et al.*, *Molecular Biology of the Cell*, 3:13-27 (1992); Ducommun *et al.*, *EMBO Journal*, 10:3311-3319 (1991); Gautier *et al.*, *Nature* 339:626-629 (1989); Gould
30 and Nurse, *Nature*, 342:39-45 (1989); Krek and Nigg, *EMBO Journal*, 10:3331-3341 (1991); Solomon *et al.*, *Cell*, 63:1013-1024 (1990)). The coordinate activation

and inactivation of different cyclin/CDK complexes is necessary for normal progression through the cell cycle (Pines, *Trends in Biochemical Sciences*, 18:195-197 (1993); Sherr, *Cell*, 73:1059-1065 (1993)). Both the critical G1-S and G2-M transitions are controlled by the activation of different cyclin/CDK activities. In G1, both cyclin D/CDK4 and cyclin E/CDK2 are thought to mediate the onset of S-phase (Matsushima *et al.*, *Molecular & Cellular Biology*, 14:2066-2076 (1994); Ohtsubo and Roberts, *Science*, 259:1908-1912 (1993); Quelle *et al.*, *Genes & Development*, 7:1559-1571 (1993); Resnitzky *et al.*, *Molecular & Cellular Biology*, 14:1669-1679 (1994)). Progression through S-phase requires the activity of cyclin A/CDK2 (Girard *et al.*, *Cell*, 67:1169-1179 (1991); Pagano *et al.*, *EMBO Journal*, 11:961-971 (1992); Rosenblatt *et al.*, *Proceedings of the National Academy of Science USA*, 89:2824-2828 (1992); Walker and Maller, *Nature*, 354:314-317 (1991); Zindy *et al.*, *Biochemical & Biophysical Research Communications*, 182:1144-1154 (1992)) whereas the activation of cyclin A/cdc2 (CDK1) and cyclin B/cdc2 are required for the onset of metaphase (Draetta, *Trends in Cell Biology*, 3:287-289 (1993)); Murray and Kirschner, *Nature*, 339:275-280 (1989); Solomon *et al.*, *Molecular Biology of the Cell*, 3:13-27 (1992); Girard *et al.*, *Cell*, 67:1169-1179 (1991); Pagano *et al.*, *EMBO Journal*, 11:961-971 (1992); Rosenblatt *et al.*, *Proceedings of the National Academy of Science USA*, 89:2824-2828 (1992); Walker and Maller, *Nature*, 354:314-317 (1991); Zindy *et al.*, *Biochemical & Biophysical Research Communications*, 182:1144-1154 (1992)). It is not surprising, therefore, that the loss of control of CDK regulation is a frequent event in hyperproliferative diseases and cancer. (Pines, *Current Opinion in Cell Biology*, 4:144-148 (1992); Lees, *Current Opinion in Cell Biology*, 7:773-780 (1995); Hunter and Pines, *Cell*, 79:573-582 (1994)). The selective inhibition of CDKs is therefore an object of the present invention.

The compounds of the present invention are additionally useful in the treatment of one or more diseases afflicting mammals which are characterized by cellular proliferation in the areas of blood vessel proliferative disorders, fibrotic disorders, mesangial cell proliferative disorders and metabolic diseases. Blood vessel proliferative disorders include arthritis and restenosis. Fibrotic disorders

include hepatic cirrhosis and atherosclerosis. Mesangial cell proliferative disorders include glomerulonephritis, diabetic nephropathy, malignant nephrosclerosis, thrombotic microangiopathy syndromes, organ transplant rejection and glomerulopathies. Metabolic disorders include psoriasis, diabetes mellitus, chronic wound healing, inflammation, neurodegenerative diseases, macular degeneration, and diabetic retinopathy.

Inhibitors of kinases involved in mediating or maintaining these disease states represent novel therapies for these disorders. Examples of such kinases include, but are not limited to: (1) inhibition of c-Src (Brickell, *Critical Reviews in Oncogenesis*, 3:401-406 (1992); Courtneidge, *Seminars in Cancer Biology*, 5:236-246 (1994), raf (Powis, *Pharmacology & Therapeutics*, 62:57-95 (1994)) and the cyclin-dependent kinases (CDKs) 1, 2 and 4 in cancer (Pines, *Current Opinion in Cell Biology*, 4:144-148 (1992); Lees, *Current Opinion in Cell Biology*, 7:773-780 (1995); Hunter and Pines, *Cell*, 79:573-582 (1994)), (2) inhibition of CDK2 or PDGF-R kinase in restenosis (Buchdunger *et al.*, *Proceedings of the National Academy of Science USA*, 92:2258-2262 (1995)), (3) inhibition of CDK5 and GSK3 kinases in Alzheimers (Hosoi *et al.*, *Journal of Biochemistry (Tokyo)*, 117:741-749 (1995); Aplin *et al.*, *Journal of Neurochemistry*, 67:699-707 (1996), (4) inhibition of c-Src kinase in osteoporosis (Tanaka *et al.*, *Nature*, 383:528-531 (1996), (5) inhibition of GSK-3 kinase in type-2 diabetes (Borthwick *et al.*, *Biochemical & Biophysical Research Communications*, 210:738-745 (1995), (6) inhibition of the p38 kinase in inflammation (Badger *et al.*, *The Journal of Pharmacology and Experimental Therapeutics*, 279:1453-1461 (1996)), (7) inhibition of VEGF-R 1-3 and TIE-1 and -2 kinases in diseases which involve angiogenesis (Shawver *et al.*, *Drug Discovery Today*, 2:50-63 (1997)), (8) inhibition of UL97 kinase in viral infections (He *et al.*, *Journal of Virology*, 71:405-411 (1997)), (9) inhibition of CSF-1R kinase in bone and hematopoietic diseases (Myers *et al.*, *Bioorganic & Medicinal Chemistry Letters*, 7:421-424 (1997), and (10) inhibition of Lck kinase in autoimmune diseases and transplant rejection (Myers *et al.*, *Bioorganic & Medicinal Chemistry Letters*, 7:417-420 (1997)).

-25-

It is additionally possible that inhibitors of certain kinases may have utility in the treatment of diseases when the kinase is not misregulated, but it nonetheless essential for maintenance of the disease state. In this case, inhibition of the kinase activity would act either as a cure or palliative for these diseases. For example,

5 many viruses, such as human papilloma virus, disrupt the cell cycle and drive cells into the S-phase of the cell cycle (Vousden, *FASEB Journal*, 7:8720879 (1993)). Preventing cells from entering DNA synthesis after viral infection by inhibition of essential S-phase initiating activities such as CDK2, may disrupt the virus life cycle by preventing virus replication. This same principle may be used to protect normal

10 cells of the body from toxicity of cycle-specific chemotherapeutic agents (Stone *et al.*, *Cancer Research*, 56:3199-3202 (1996); Kohn *et al.*, *Journal of Cellular Biochemistry*, 54:44-452 (1994)). Inhibition of CDKs 2 or 4 will prevent progression into the cycle in normal cells and limit the toxicity of cytotoxics which act in S-phase, G2 or mitosis. Furthermore, CDK2/cyclin E activity has also been

15 shown to regulate NF-kB. Inhibition of CDK2 activity stimulates NF-kB-dependent gene expression, an event mediated through interactions with the p300 coactivator (Perkins *et al.*, *Science*, 275:523-527 (1997)). NF-kB regulates genes involved in inflammatory responses (such as hematopoietic growth factors, chemokines and leukocyte adhesion molecules) (Baeuerle and Henkel, *Annual Review of*

20 *Immunology*, 12:141-179 (1994)) and may be involved in the suppression of apoptotic signals within the cell (Beg and Baltimore, *Science*, 274:782-784 (1996); Wang *et al.*, *Science*, 274:784-787 (1996); Van Antwerp *et al.*, *Science*, 274:787-789 (1996)). Thus, inhibition of CDK2 may suppress apoptosis induced by cytotoxic drugs via a mechanism which involves NF-kB. This therefore suggests that

25 inhibition of CDK2 activity may also have utility in other cases where regulation of NF-kB plays a role in etiology of disease. A further example may be take from fungal infections: Aspergillosis is a common infection in immune-compromised patients (Armstrong, *Clinical Infectious Diseases*, 16:1-7 (1993)). Inhibition of the Aspergillus kinases Cdc2/CDC28 or Nim A (Osmani *et al.*, *EMBO Journal*,

30 10:2669-2679 (1991); Osmani *et al.*, *Cell*, 67:283-291 (1991)) may cause arrest or death in the fungi, improving the therapeutic outcome for patients with these infections.

-26-

In one embodiment, the present invention provides compounds of formulas 1-109 as described above. The values of substituents in preferred groups of compounds of formulas 1-109 are given below.

Preferably, R_b is selected from the group consisting of F, Cl, Br, I, CH_3 , NO_2 , OCF_3 , OCH_3 , CN, CO_2CH_3 , CF_3 , t-butyl, pyridyl, substituted or unsubstituted oxazolyl, substituted or unsubstituted benzyl, substituted or unsubstituted benzenesulfonyl, substituted or unsubstituted phenoxy, substituted or unsubstituted phenyl, substituted or unsubstituted amino, carboxyl, substituted and unsubstituted tetrazolyl, substituted and unsubstituted styryl, substituted and unsubstituted arylthio, substituted and unsubstituted heteroarylthio; CH_2OR_c , wherein R_c is hydrogen or substituted or unsubstituted alkyl or aryl; and $-W-(CH_2)_t-NR_dR_e$, wherein t is an integer from about 1 to about 6; W is a direct bond, O, S, $S(O)$, $S(O)_2$, or NR_f , wherein R_f is H or alkyl and R_d and R_e are independently H, alkyl, alkanoyl or SO_2 -alkyl; or R_d , R_e and the nitrogen atom to which they are attached together form a five- or six-membered heterocyclic ring.

Preferably R_a is selected from the group consisting of F, Cl, Br, I, CH_3 , NO_2 , OCF_3 , OCH_3 , CN, CO_2CH_3 , CF_3 , t-butyl, pyridyl, substituted or unsubstituted oxazolyl, substituted or unsubstituted benzyl, substituted or unsubstituted benzenesulfonyl, substituted or unsubstituted phenoxy, substituted or unsubstituted phenyl, substituted or unsubstituted amino, carboxyl, substituted and unsubstituted tetrazolyl, substituted and unsubstituted styryl, substituted and unsubstituted arylthio, substituted and unsubstituted heteroarylthio; CH_2OR_c , wherein R_c is hydrogen or substituted or unsubstituted alkyl or aryl; and $-W-(CH_2)_t-NR_dR_e$, wherein t is an integer from about 1 to about 6; W is a direct bond, O, S, $S(O)$, $S(O)_2$, or NR_f , wherein R_f is H or alkyl and R_d and R_e are independently H, alkyl, alkanoyl or SO_2 -alkyl; or R_d , R_e and the nitrogen atom to which they are attached together form a five- or six-membered heterocyclic ring.

In one embodiment, R_2 is an oxacycloalkyl group of the formula




-27-

wherein n is 1, 2 or 3.

In another embodiment, R_2 is of the formula

5




where m is 0, 1, 2 or 3 and R_9 is H or $-(CH_2)_pN(R_4)R_5$, where p is an integer from about 2 to about 6. R_4 and R_5 are each, independently, H, azabicycloalkyl or Y-Z, wherein Y is selected from the group consisting of $-C(O)-$, $-(CH_2)_p-$, $-S(O)_2-$, $-C(O)O-$, $-SO_2NH-$, $-CONH-$, $(CH_2)_qO-$, $-(CH_2)_qNH-$, and $-(CH_2)_qS(O)_r-$; wherein p is an integer from 0 to about 6, q is an integer from 0 to about 6, and r is 0, 1 or 2; and Z is a substituted or unsubstituted alkyl, amino, aryl, heteroaryl or heterocycloalkyl group or R_4 , R_5 and the nitrogen atom together form a 3, 4, 5, 6 or 7-membered, substituted or unsubstituted heterocyclic or heterobicyclic group.

15

In another embodiment, R_2 is of the formula

20



wherein m is 1, 2 or 3. a and b are each, independently, an integer from 0 to about, except that when the two substituents are attached to the same carbon atom, a is from 1 to about 6. Q is NR_4R_5 or $-OR_6$. Each R_4 and R_5 is, independently, H, azabicycloalkyl or Y-Z, wherein Y is selected from the group consisting of $-C(O)-$, $-(CH_2)_p-$, $-S(O)_2-$, $-C(O)O-$, $-SO_2NH-$, $-CONH-$, $(CH_2)_qO-$, $-(CH_2)_qNH-$, and $-(CH_2)_qS(O)_r-$; where p is an integer from 0 to about 6, q is an integer from 0 to about 6, and r is 0, 1 or 2; and Z is a substituted or unsubstituted alkyl, amino, aryl, heteroaryl or heterocycloalkyl group. R_4 , R_5 and the nitrogen atom can also together form a 3, 4, 5, 6 or 7-membered, substituted or unsubstituted heterocyclic or heterobicyclic group.


25

30

-28-

In another embodiment, R_2 is of the formula

5




where n is 1, 2 or 3; and R_4 is H, azabicycloalkyl or Y-Z, wherein Y is selected from the group consisting of $-C(O)-$, $-(CH_2)_p-$, $-S(O)_2-$, $-C(O)O-$, $-SO_2NH-$, $-CONH-$, $(CH_2)_qO-$, $-(CH_2)_qNH-$, and $-(CH_2)_qS(O)_r-$; wherein p is an integer from 0 to about 6, q is an integer from 0 to about 6, and r is 0, 1 or 2; and Z is a substituted or unsubstituted alkyl, amino, aryl, heteroaryl or heterocycloalkyl group.

10

In another embodiment, R_2 is of the formula

15



where m is 0, 1, 2 or 3. R_5 is H, azabicycloalkyl or Y-Z, where Y is selected from the group consisting of $-C(O)-$, $-(CH_2)_p-$, $-S(O)_2-$, $-C(O)O-$, $-SO_2NH-$, $-CONH-$, $-(CH_2)_qO-$, $-(CH_2)_qNH-$, and $-(CH_2)_qS(O)_r-$; where p is an integer from 0 to about 6, q is an integer from 0 to about 6, and r is 0, 1 or 2; and Z is a substituted or unsubstituted alkyl, amino, aryl, heteroaryl or heterocycloalkyl group. R_6 represents one or more substituents independently selected from the group consisting of hydrogen, hydroxy, oxo and substituted or unsubstituted alkyl, aryl, heteroaryl, alkoxy, carbonyl, amino, and arylalkyl groups, provided that the carbon atoms adjacent to the nitrogen atom are not substituted by a hydroxy group.


20

25

-29-

In another embodiment, R_2 is of the formula


5



- 10 wherein R_4 is H, azabicycloalkyl or Y-Z, wherein Y is selected from the group consisting of $-C(O)-$, $-(CH_2)_p-$, $-S(O)_2-$, $-C(O)O-$, $-SO_2NH-$, $-CONH-$, $(CH_2)_qO-$, $-(CH_2)_qNH-$, and $-(CH_2)_qS(O)_r-$; wherein p is an integer from 0 to about 6, q is an integer from 0 to about 6, and r is 0, 1 or 2; and Z is a substituted or unsubstituted alkyl, amino, aryl, heteroaryl or heterocycloalkyl group.

- 15 In another embodiment, R_2 is of the formula


20



- 25 where m is an integer from 1 to about 6; and R_4 and R_5 are each, independently, H, azabicycloalkyl or Y-Z, wherein Y is selected from the group consisting of $-C(O)-$, $(CH_2)_p-$, $-S(O)_2-$, $-C(O)O-$, $-SO_2NH-$, $-CONH-$, $(CH_2)_qO-$, $-(CH_2)_qNH-$, and $-(CH_2)_qS(O)_r-$; wherein p is an integer from 0 to about 6, q is an integer from 0 to about 6, and r is 0, 1 or 2; and Z is a substituted or unsubstituted alkyl, amino, aryl, heteroaryl or heterocycloalkyl group. R_4 , R_5 and the nitrogen atom can also together form a 3, 4, 5, 6 or 7-membered, substituted or unsubstituted heterocyclic or heterobicyclic group.

-30-

In another embodiment, R_2 is of the formula




where n is an integer from 0 to about 4; and r is 0 or 1. When r is 0, m is an integer from 0 to 6. When r is 1, m is an integer from 1 to 6. Q is $-NR_4R_5$ or $-OR_6$. Each R_4 and R_5 is, independently, H, azabicycloalkyl or Y-Z, wherein Y is selected from the group consisting of $-C(O)-$, $-(CH_2)_p-$, $-S(O)_2-$, $-C(O)O-$, $-SO_2NH-$, $-CONH-$, $(CH_2)_qO-$, $-(CH_2)_qNH-$, and $-(CH_2)_qS(O)_r-$; wherein p is an integer from 0 to about 6, q is an integer from 0 to about 6, and r is 0, 1 or 2; and Z is a substituted or unsubstituted alkyl, amino, aryl, heteroaryl or heterocycloalkyl group. R_4 , R_5 and the nitrogen atom can also together form a 3, 4, 5 or 6-membered, substituted or unsubstituted heterocyclic group. R_6 is hydrogen or a substituted or unsubstituted alkyl group.

15

20

In another embodiment, R_2 is of the formula




where n is an integer from 0 to about 4 and m is an integer from 0 to about 6. R_4 is H, azabicycloalkyl or Y-Z, wherein Y is selected from the group consisting of -

-31-


C(O)-, $-(CH_2)_p-$, $-S(O)_2-$, $-C(O)O-$, $-SO_2NH-$, $-CONH-$, $(CH_2)_qO-$, $-(CH_2)_qNH-$, and $(CH_2)_qS(O)_r-$; wherein p is an integer from 0 to about 6, q is an integer from 0 to about 6, and r is 0, 1 or 2; and Z is a substituted or unsubstituted alkyl, amino, aryl, heteroaryl or heterocycloalkyl group. R_6 is hydrogen or a substituted or unsubstituted alkyl group.

In embodiments of R_2 described above which include an $-N(R_4)R_5$ group, this group can form a heterocyclic group of the formula



where R_7 , R_8 , R_9 , R_{10} , R_{11} , R_{12} , R_{13} and R_{14} are each, independently, lower alkyl or hydrogen; or at least one pair of substituents R_7 and R_8 ; R_9 and R_{10} ; R_{11} and R_{12} ; or R_{13} and R_{14} together are an oxygen atom; or at least one of R_7 and R_9 is cyano, $CONHR_{15}$, $COOR_{15}$, CH_2OR_{15} or $CH_2NR_{15}(R_{16})$, where R_{15} and R_{16} are each, independently, H, azabicycloalkyl or Y-Z, wherein Y is selected from the group consisting of $-C(O)-$, $-(CH_2)_p-$, $-S(O)_2-$, $-C(O)O-$, $-SO_2NH-$, $-CONH-$, $(CH_2)_qO-$, $-(CH_2)_qNH-$, and $-(CH_2)_qS(O)_r-$; wherein p is an integer from 0 to about 6, q is an integer from 0 to about 6, and r is 0, 1 or 2; and Z is a substituted or unsubstituted alkyl, amino, aryl, heteroaryl or heterocycloalkyl group; or R_{15} , R_{16} and the nitrogen atom together form a 3, 4, 5, 6 or 7-membered, substituted or unsubstituted heterocyclic or heterobicyclic group; X is O, S, SO, SO_2 , CH_2 , $CHOR_{17}$ or NR_{17} , wherein R_{17} is hydrogen, substituted or unsubstituted alkyl, aryl, arylalkyl, $-C(NH)NH_2$, $-C(O)R_{17}$, or $-C(O)OR_{18}$, wherein R_{18} is hydrogen, substituted or unsubstituted alkyl, aryl or arylalkyl; and t is 0 or 1.

R_4 , R_5 and the nitrogen atom can also together form a heterocyclic group of the formula



-32-

where R_{19} and R_{20} are each, independently, hydrogen or lower alkyl; or R_{19} and R_{20} together are an oxygen atom. R_{21} and R_{22} are each, independently, H azabicycloalkyl or Y-Z, wherein Y is selected from the group consisting of -C(O)-, $-(CH_2)_p-$, $-S(O)_2-$, -C(O)O-, $-SO_2NH-$, $-CONH-$, $(CH_2)_qO-$, $-(CH_2)_qNH-$, and $-(CH_2)_qS(O)_r-$; wherein p is an integer from 0 to about 6, q is an integer from 0 to about 6, and r is 0, 1 or 2, and Z is a substituted or unsubstituted alkyl, amino, aryl, heteroaryl or heterocycloalkyl group. R_{21} , R_{22} and the nitrogen atom can also together form a 3, 4, 5 or 6-membered, substituted or unsubstituted heterocyclic group. m is an integer from 1 to about 6; and n is an integer from 0 to about 6.

R_4 , R_5 and the nitrogen atom can also together form a heterocyclic group of the formula



where m is an integer from 1 to 6. R_{23} is CH_2OH , NRR' , $C(O)NR'R$ or $COOR$, where R and R' are each independently hydrogen or a substituted or unsubstituted alkyl, aryl or arylalkyl group.

R_4 , R_5 and the nitrogen atom can also together form a heterocyclic group of the formula




where R_{24} is a substituted or unsubstituted alkyl, aryl or arylalkyl group, carboxyl, cyano, $C(O)OR_{25}$, CH_2OR_{25} , $CH_2NR_{26}R_{27}$ or $C(O)NHR_{26}$. R_{25} is a substituted or unsubstituted alkyl, aryl, arylalkyl, heterocyclic or heteroaryl group. R_{26} and R_{27} are each, independently, H, azabicycloalkyl or Y-Z, wherein Y is selected from the group consisting of -C(O)-, $-(CH_2)_p-$, $-S(O)_2-$, -C(O)O-, $-SO_2NH-$, $-CONH-$, $(CH_2)_qO-$

-(CH₂)_qNH-, and -(CH₂)_qS(O)_r-; wherein p is an integer from 0 to about 6, q is an integer from 0 to about 6, and r is 0, 1 or 2; and Z is a substituted or unsubstituted alkyl, amino, aryl, heteroaryl or heterocycloalkyl group. R₂₆, R₂₇, and the nitrogen atom can also together form a 3, 4, 5 or 6-membered, substituted or unsubstituted heterocyclic group.

In one subset of compounds of Formulas 1-109, at least one of R₄ and R₅ is of the formula Y-Z, where Z is of the formula

10



where T is C(O), S, SO, SO₂, CHOR or NR, wherein R is hydrogen or a substituted or unsubstituted alkyl, aryl or arylalkyl group; and n is 0, 1 or 2.


In another embodiment, at least one of R₄ and R₅ is of the formula Y-Z, where Z is -N(R₂₈)R₂₉, and R₂₈ and R₂₉ are each, independently, substituted or unsubstituted carboxyalkyl, alkoxycarbonylalkyl, hydroxyalkyl, alkylsulfonyl, alkylcarbonyl or cyanoalkyl. R₂₈ and R₂₉, together with the nitrogen atom, can also form a five- or six-membered heterocyclic group.

In yet another embodiment, at least one of R₄ and R₅ is of the formula Y-Z, where Z is of

the formula N(R₃₀)R₃₁. R₃₀ and R₃₁ are each, independently, hydrogen, alkyl, alkoxycarbonyl, alkoxyalkyl, hydroxyalkyl, aminocarbonyl, cyano, alkylcarbonyl or arylalkyl.

In another embodiment, at least one of R₄ and R₅ is Y-Z, where Z is of the formula

30




Each X is, independently, CH or N. R₃₂ is hydrogen, cyano or a substituted or unsubstituted alkyl, alkoxycarbonyl, alkoxyalkyl, hydroxyalkyl, aminocarbonyl,

-34-

alkylcarbonyl or arylalkyl group.

One of R_4 and R_5 can also be Y-Z where Z is of the formula



where g is 0 or 1; and T is $C(O)$, O , S , SO , SO_2 , CH_2 , $CHOR_{17}$ or NR_{17} . R_{17} is
 10 hydrogen, substituted or unsubstituted alkyl, aryl, arylalkyl, $-C(NH)NH_2$, $-C(O)R_{18}$,
 $C(O)NH_2$ or $-C(O)OR_{18}$, where R_{18} is hydrogen, substituted or unsubstituted alkyl,
 aryl or arylalkyl. R_{32} is hydrogen, cyano or a substituted or unsubstituted alkyl,
 alkoxycarbonyl, alkoxyalkyl, hydroxyalkyl, aminocarbonyl, alkylcarbonyl or
 arylalkyl group.

15 One of R_4 and R_5 can also be Y-Z, where Z is of the formula



where g is 0, 1 or 2; and R_{32} is hydrogen, cyano or a substituted or unsubstituted
 alkyl, alkoxycarbonyl, alkoxyalkyl, hydroxyalkyl, aminocarbonyl, alkylcarbonyl or
 arylalkyl group.

Z can also be of the formula




where g is 0, 1, 2 or 3, and T is O , S , SO , SO_2 , CH_2 , $CHOR_{17}$ or NR_{17} . R_{17} is
 30 hydrogen, substituted or unsubstituted alkyl, aryl, arylalkyl, $-C(NH)NH_2$, $-C(O)R_{17}$,
 or $-C(O)OR_{18}$, wherein R_{18} is hydrogen, substituted or unsubstituted alkyl, aryl or
 arylalkyl. R_{32} is hydrogen, cyano or a substituted or unsubstituted alkyl,

-35-


alkoxycarbonyl, alkoxyalkyl, hydroxyalkyl, aminocarbonyl, alkylcarbonyl or arylalkyl group.

One of R_4 and R_5 can also be Y-Z, wherein Z is of the formula



Where R_{32} is hydrogen, cyano or substituted or unsubstituted alkyl, alkoxycarbonyl, alkoxyalkyl, hydroxyalkyl, aminocarbonyl, alkylcarbonyl, thioalkoxy or arylalkyl; and R_{33} is hydrogen or substituted or unsubstituted alkyl, alkoxycarbonyl, alkoxyalkyl, aminocarbonyl, perhaloalkyl, alkenyl, alkylcarbonyl or arylalkyl.


In another subset of the compounds of Formulas 1-109, R_2 is of the formula



where m is 0 or 1; R_{34} , R_{35} , R_{36} , R_{37} , R_{38} , R_{39} , R_{40} and R_{41} are each, independently, methyl or hydrogen; or at least one pair of substituents R_{34} and R_{35} ; R_{36} and R_{37} ; R_{38} and R_{39} ; or R_{40} and R_{41} together are an oxygen atom. R_{42} is H, azabicycloalkyl or Y-Z, wherein Y is selected from the group consisting of $-C(O)-$, $-(CH_2)_p-$, $-S(O)_2-$, $-C(O)O-$, $-SO_2NH-$, $-CONH-$, $(CH_2)_qO-$, $-(CH_2)_qNH-$, and $-(CH_2)_qS(O)_r-$; wherein p is an integer from 0 to about 6, q is an integer from 0 to about 6, and r is 0, 1 or 2; and Z is a substituted or unsubstituted alkyl, amino, aryl, heteroaryl or heterocycloalkyl group.


-36-

In a preferred embodiment, R_{42} is of the formula



Where u is 0 or 1; R_{43} , R_{44} , R_{45} , R_{46} , R_{47} , R_{48} , R_{49} and R_{50} are each, independently, methyl or hydrogen; or at least one pair of substituents R_{43} and R_{44} ; R_{45} and R_{46} ; R_{47} and R_{48} ; or R_{49} and R_{50} together are an oxygen atom. R_{51} is H, azabicycloalkyl or V-L, where V is selected from the group consisting of $-C(O)-$, $-(CH_2)_p-$, $-S(O)_2-$, $-C(O)O-$, $-SO_2NH-$, $-CONH-$, $(CH_2)_qO-$, $-(CH_2)_qNH-$, and $-(CH_2)_qS(O)_r-$; wherein p is an integer from 0 to about 6, q is an integer from 0 to about 6, and r is 0, 1 or 2; and L is a substituted or unsubstituted alkyl, amino, aryl, heteroaryl or heterocycloalkyl group.


In another subset of the compounds of Formulas 1-109, R_2 is of the formula



Where h , i , j , k and l are independently 0 or 1; R_{52} , R_{53} , R_{54} , R_{55} , R_{56} , R_{57} , R_{58} , R_{59} , R_g and R_h are each, independently, methyl or hydrogen; or at least one pair of substituents R_{52} and R_{53} ; R_{54} and R_{55} ; R_{56} and R_{57} ; or R_{58} and R_{59} together are an oxygen atom. R_{60} is H, azabicycloalkyl or Y-Z, wherein Y is selected from the group consisting of $-C(O)-$, $-(CH_2)_p-$, $-S(O)_2-$, $-C(O)O-$, $-SO_2NH-$, $-CONH-$, $(CH_2)_qO-$, $-(CH_2)_qNH-$, and $-(CH_2)_qS(O)_r-$; wherein p is an integer from 0 to about 6, q is an integer from 0 to about 6, and r is 0, 1 or 2; and Z is a substituted or unsubstituted

-37-

alkyl, amino, aryl, heteroaryl or heterocycloalkyl group. In one embodiment, R_{60} is of the formula



Where v is 0 or 1; R_{61} , R_{62} , R_{63} , R_{64} , R_{65} , R_{66} , R_{67} and R_{68} are each, independently, lower alkyl or hydrogen; or at least one pair of substituents R_{61} and R_{62} ; R_{63} and R_{64} ; R_{65} and R_{66} ; and R_{67} and R_{68} together are an oxygen atom; and R_{69} is H, azabicycloalkyl or V-L, where V is selected from the group consisting of $-\text{C}(\text{O})-$, $-(\text{CH}_2)_p-$, $-\text{S}(\text{O})_2-$, $-\text{C}(\text{O})\text{O}-$, $-\text{SO}_2\text{NH}-$, $-\text{CONH}-$, $(\text{CH}_2)_q\text{O}-$, $-(\text{CH}_2)_q\text{NH}-$, and $(\text{CH}_2)_q\text{S}(\text{O})_r-$; wherein p is an integer from 0 to about 6, q is an integer from 0 to about 6, and r is 0, 1 or 2; and L is a substituted or unsubstituted alkyl, amino, aryl, heteroaryl or heterocycloalkyl group.


In another subset of compounds of Formula (I), R_3 is H;

R_2 is $-\text{Z}^{101}-\text{Z}^{102}$ where Z^{101} is a covalent bond, $-(\text{C}_1-\text{C}_6)-$, $-(\text{C}_1-\text{C}_6)-\text{O}-$, $-(\text{C}_1-\text{C}_6)-\text{C}(\text{O})-$, $-(\text{C}_1-\text{C}_6)-\text{C}(\text{O})\text{O}-$, $-(\text{C}_1-\text{C}_6)-\text{C}(\text{O})-\text{NH}-$, $-(\text{C}_1-\text{C}_6)-\text{C}(\text{O})-\text{N}((\text{C}_1-\text{C}_6))-$ or a substituted phenyl group; and

Z^{102} is hydrogen, a substituted or unsubstituted alkyl group or a substituted or unsubstituted, saturated or unsaturated heterocyclic group.


In another subset of compounds of Formula (I) Z^{101} is selected from the group consisting of $-\text{CH}_2-\text{C}(\text{O})\text{O}-$, $-\text{CH}_2-\text{C}(\text{O})-$, $-\text{CH}_2-\text{C}(\text{O})-\text{NH}-$, $-\text{CH}_2-\text{C}(\text{O})-\text{N}(\text{Me})-$, $-\text{CH}(\text{Me})-\text{C}(\text{O})\text{O}-$, $-(\text{CH}_2)_3-\text{C}(\text{O})\text{O}-$, $-\text{CH}(\text{Me})-\text{C}(\text{O})-\text{NH}-$, and $-(\text{CH}_2)_3-\text{C}(\text{O})-\text{NH}-$; Z^{102} is selected from the group consisting of hydrogen, methyl, ethyl, N,N-dimethylaminoethyl, N,N-diethylaminoethyl, 2-phenyl-2-hydroxyethyl, morpholino, piperazinyl, N-methylpiperazinyl and 2-hydroxymethylpyrrolidinyl.

-38-



a substituted or unsubstituted benzoxazolyl or a substituted or unsubstituted benzthiazolyl.


5 In another subset of compounds of Formula (I), R₁ is




where there is only one R_a and it is H or F.

10 In another subset of compounds of Formula (I) Z¹⁰¹ is a covalent bond; and Z¹⁰² is an optionally substituted pyridyl.

In another subset of compounds of Formula (I), R₁ is



In another subset of compounds of Formula (I), R₃ is H;
R₂ is cyclopentyl; and



In another subset of compounds of Formula (I), Z^{110} is hydrogen;
A is O; and Z^{100} is optionally substituted phenyl, furanyl or thienyl, where Z^{100} is optionally substituted with one or more substituents each independently selected from the group consisting of F, COOH, NO₂, OMe, -COOMe, OCF₃ and CF₃.

5 In another subset of compounds of Formula (I), Z^{110} is hydrogen;

A is -O-, -O-(CR₂)_n-C(O)- or -O-(CR₂)_n-O-;

n for each occurrence is 0 to 3;

Z^{100} is an optionally substituted group selected from the group consisting of cyclohexyl, phenyl, tetrahydropyranyl, tetrahydrofuranyl, isoxazolyl and piperidinyl;
10 where Z^{100} is optionally substituted with one or more substituents selected from the group consisting of alkyl, alkoxy, halo, hydroxy and alkoxy carbonyl.

In another subset of compounds of Formula (I), R² is an optionally substituted group selected from the group consisting of cyclobutyl and cyclohexyl.

In another subset of compounds of Formula (I), R² is optionally substituted
15 with one or more substituents selected from the group consisting of hydroxy, alkyl, hydroxyalkyl, carboxyalkyl and phenylalkoxyalkyl.

In another subset of compounds of Formula (I), R₁ is 4-phenoxyphenyl.

In another subset of compounds of Formula (I), m is 2; a is 0; R₆ is H; b is 1 or 2; and R₄ and R₅ are each hydrogen.

20 In another subset of compounds of Formula (I), m is 0, 1 or 2; R₆ is hydrogen; R₅ is H or Y-Z;


where Y is a covalent bond, -C(O)-, -(CH₂)_qO-, -(CH₂)_q-, -(CH₂)_qC(O)- or -C(O)(CH₂)_q-, where the alkyl portion of -(CH₂)_qO-, -(CH₂)_p-, -(CH₂)_qC(O)- and -C(O)(CH₂)_q- is optionally substituted by a halogen, hydroxy or an alkyl
25 group; and

Z is hydrogen, alkyl, optionally substituted alkyl, alkoxyalkyl, optionally substituted heterocycloalkyl, optionally substituted heteroaryl, or optionally substituted amino.


In another subset of compounds of Formula (I), Z is hydrogen, methyl, ethyl,
30 hydroxymethyl, methoxyethyl, N-methyl-piperidinyl, (t-butoxycarbonyl)(hydroxy)-piperidinyl, hydroxypiperidinyl, (hydroxymethyl)piperidinyl, (hydroxy)(methyl)-

-40-


- 5 piperidinyl, morpholino, (methoxyethyl)piperiziny, methylpiperiziny, 4-piperidinylpiperidinyl, imidazolyl, methylimidazolyl, N-methylamino, N,N-dimethylamino, N-isopropylamino, N,N-diethylamino, 2,3-dihydroxypropylamino, 2-hydroxyethylamino, 3-hydroxypropylamino, methoxyethylamino, ethoxycarbonylmethylamino, phenylmethylamino, N-methyl-N-methoxyamino,



- , furanylmethylamino, piperidinyethylamino, N-(2-N,N-dimethylaminoethyl)-N-methylamino, 2-N,N-dimethylaminoethylamino, N-methyl-N-(N-methylpiperidin-4-yl)amino, 2-morpholino-ethylamino, 3-morpholino-propylamino, 3-imidazolylpropylamino, or 3-(2-oxopyrrolidinyl)propylamino.
- 10 In another subset of compounds of Formula (I), m is 2; R₅ is Y-Z; Y is -



In another subset of compounds of Formula (I), R₄ is hydrogen or methyl;



- A is selected from the group consisting of O, -N(R)- and -N(R)C(O)-;
- 15 Z¹¹¹ is -(CH₂)_n-cycloalkyl-(CH₂)_n-;

R is hydrogen or alkyl;

n is 0 to 5;

R_a is one or more substituents each independently selected from the group consisting of H, OH, F, Cl, methyl and methoxy; and

- 20 R_b is one or more substituents each independently selected from the group consisting of H, CN, F, CF₃, OCF₃, methyl, methoxy and an optionally substituted amino group;


where said amino group is optionally substituted with one or two groups each independently selected from the group consisting of

-41-

alkyl, alkoxyalkyl, phenyl, substituted phenyl, and optionally substituted heteroaryl.

In another subset of compounds of Formula (I), R_b is 4-methylphenylthio or 2-pyridinylthio.

5 In another subset of compounds of Formula (I),




R_1 is

where Z^{100} is selected from the group consisting of benzo[b]thiophene, furanyl and thiophene.




10 In another subset of compounds of Formula (I), wherein R_a is alkoxy; A is -NH-C(O)-; and there is a covalent bond between A and Z^{100} .

In another subset of compounds of Formula (I),



R_1 is

A is selected from the group consisting of -N(R)-C(O)-N(R)-, -(CH₂)_n-N(R)C(O)N(R)-, -N(R)- and -N(R)-SO₂-; R is hydrogen or alkyl;

15 Z^{100} is , , , pyridinyl, thiazolyl, furanyl, benzofuranyl or oxazolyl;

X is S, O or NR where R for each occurrence is independently H or Me;

20 R_a is one or more substituents each independently selected from the group consisting of H and F; and


R_b is one or more substituents each independently selected from the group consisting of H, F, Cl, Br, NO₂, CF₃, alkyl, alkoxy and alkoxycarbonyl.

In another subset of compounds of Formula (I), R_4 is methyl; m is 1, 2 or 3; R_5 is Y-Z, where Y is -C(O)O-, -C(O)- or -C(O)-(CH₂)_p-; and Z is aminoalkyl, N-

-42-

alkylamino, N,N-dialkylamino or hydroxyalkylaminoalkyl.

In another subset of compounds of Formula (I), R_4 is methyl; R_1 is




where n is 0 to 3; Z^{100} is an optionally

substituted group selected from the group consisting of indolyl, indenyl,

5 methylindenyl, methylindolyl, dimethylaminophenyl, phenyl, cyclohexyl and benzofuranyl.

In another subset of compounds of Formula (I),



Z^{100} is an optionally substituted group selected from the group consisting of

10 phenyl, imidazolyl, indolyl, furanyl, benzofuranyl and 2,3-dihydrobenzofuranyl;

where Z^{100} is optionally substituted with one or more substituents
each independently selected from the group consisting of F, Cl, CN,
optionally substituted alkyl, -O-(optionally substituted alkyl), -
COOH, $-Z^{105}-C(O)N(R)_2$, $-Z^{105}-N(R)-C(O)-Z^{200}$, $-Z^{105}-N(R)-S(O)_2-$
15 Z^{200} , and $-Z^{105}-N(R)-C(O)-N(R)-Z^{200}$;

Z^{105} is a covalent bond or (C_1-C_6) ;

Z^{200} is an optionally substituted group selected from group
consisting of (C_1-C_6) , phenyl and $-(C_1-C_6)$ -phenyl;

Z^{110} and Z^{111} are each independently a covalent bond or (C_1-C_3) group optionally


20 substituted with alkyl, hydroxy, COOH, CN or phenyl; and

A is O, $-N(R)-C(O)-N(R)-$, $-N(R)-C(O)-O-$, $-N(R)-$ or $-N(R)-C(O)-$, where R is
H or alkyl.

In another subset of compounds of Formula (I), R_4 is methyl.

In another subset of compounds of Formula (I),


-43-




where Z^{100} is an optionally substituted group selected from the group consisting of benzoxazolyl, benzothiazolyl and benzimidazolyl.

- In another subset of compounds of Formula (I), R_4 is methyl; A is -NH-;
 5 there is only one R_a and it is H or F; and Z^{100} is optionally substituted with one or more substituents each independently selected from the group consisting of alkyl, halo, CF_3 , and alkoxy.

In another subset of compounds of Formula (I),



- 10 Z^{100} is an optionally substituted group selected from the group consisting of phenyl, pyrrolyl, pyridyl, benzimidazolyl, naphthyl and



- 15 where Z^{100} is optionally substituted with one or more substituents each independently selected from the group consisting of F, Cl, Br, NO_2 , amino, N-alkylamino, N,N-dialkylamino, CN, optionally substituted alkyl, -O-(optionally substituted alkyl) and phenyl;


Z^{110} and Z^{111} for each occurrence is independently (C_0 - C_3) optionally substituted with optionally substituted phenyl; and

A is -N(R)-C(O)-N(R)-, -N(R)-S(O)₂-, -N(R)-C(O)-, -N(R)- or -N(R)-C(O)-O-.

- 20 In another subset of compounds of Formula (I), R_4 is methyl and there is only one R_a and it is F.

In another subset of compounds of Formula (I),

-44-



Z^{100} is an optionally substituted group selected from the group consisting of phenyl, isoxazolyl, tetrahydronaphthyl, furanyl, benzofuranyl, pyridyl and indolyl;

5 where Z^{100} is optionally substituted with one or more substituents each independently selected from the group consisting of F, CN, NO_2 , -C(O)H, -CONH₂, -NHSO₂CF₃, optionally substituted alkyl, optionally substituted heteroaryl and -O-(optionally substituted alkyl);


Z^{110} and Z^{111} are each independently optionally substituted (C_0 - C_3); and

10 A is O, -N(R)-C(O)-(CH₂)_n-N(R)-, -C(O)-N(R)-, -N(R)-C(O)-O-, -N(R)-C(O)- or -N(R)-.

In another subset of compounds of Formula (I), R_4 is methyl; R_5 is H or methoxy; and Z^{110} and Z^{111} are each unsubstituted.


In another subset of compounds of Formula (I), R_1 is

-45-



where R is H or lower alkyl and n is for each occurrence is independently 1 to 6.


In another subset of compounds of Formula (I), R_1 is



5

In another subset of compounds of Formula (I), Z^{100} is substituted or unsubstituted phenyl.

In another subset of compounds of Formula (I),



where Z^{100} is an optionally substituted group selected

-46-

























from the group consisting of benzoxazolyl, benzothiazolyl and benzimidazolyl.

In another subset of compounds of Formula (I), n is 2; R₆ is H; m is 1; r is 1; and R₄ and R₅ are each hydrogen.

In another subset of compounds of Formula (I), R₁ is 4-phenoxyphenyl.

























- 5 A subset of compounds of formulas 1-117 have R₁ = 4-phenoxyphenyl, R₂ = cyclopentyl and both R₃ = H. These compounds are illustrated below.

-47-

























 -38.36	 -38.30	 -37.83	 -36.11
 -37.84	 -37.02	 -37.05	 -36.84
 -34.22	 -33.88	 -30.20	 -32.12
 -35.04	 -39.18	 -33.11	 -31.32
 -36.74	 -36.42	 -36.33	 -37.66
 -37.70	 -38.61	 -37.77	 -32.93

-48-














 -30.27	 -30.69	 -31.88	 -30.33
 -31.67	 -31.15	 -31.44	 -31.62
 -35.14	 -30.85	 -34.81	 -34.81
 -35.08	 -31.06	 -35.35	 -33.82
 -31.81	 -33.35	 -31.29	 -28.20
 -32.39	 -32.10	 -31.66	 -31.33

 -26.77	 -28.81	 -28.48	 -27.85
 -29.87	 -29.89	 -24.06	 -26.18
 -26.70	 -28.81	 -28.51	 -27.20
 -28.06	 -24.31	 -28.76	 -27.71
 -32.01	 -28.04	 -30.32	 -27.51
 -30.14	 -31.45	 -30.58	 -28.01

-50-

 -28.83	 -28.49	 -28.35	 -28.62
 -27.00	 -34.13	 -28.69	 -31.83
 -28.79	 -32.33	 -28.17	 -32.70
 -32.99	 -32.83	 -32.16	 -29.09
 -28.25	 -32.03	 -31.73	 -31.73
 -32.71	 -29.16	 -32.77	 -29.83

-51-

 -24.20	 -24.48	
 -27.85	 -24.82	
 -25.48	 -24.07	
 -23.04	 -28.22	
 -28.99	 -27.35	
 -22.15	 -27.48	 -22.00

-52-

Compounds of formulas 1-109 may exist as salts with pharmaceutically acceptable acids. The present invention includes such salts. Examples of such salts include hydrochlorides, hydrobromides, sulfates, methanesulfonates, nitrates, maleates, acetates, citrates, fumarates, tartrates [eg (+)-tartrates, (-)-tartrates or mixtures thereof including racemic mixtures], succinates, benzoates and salts with amino acids such as glutamic acid. These salts may be prepared by methods known to those skilled in the art.

Certain compounds of formulas 1-109 which have acidic substituents may exist as salts with pharmaceutically acceptable bases. The present invention includes such salts. Example of such salts include sodium salts, potassium salts, lysine salts and arginine salts. These salts may be prepared by methods known to those skilled in the art.

Certain compounds of formulas 1-109 and their salts may exist in more than one crystal form and the present invention includes each crystal form and mixtures thereof.

Certain compounds of formulas 1-109 and their salts may also exist in the form of solvates, for example hydrates, and the present invention includes each solvate and mixtures thereof.

Certain compounds of formulas 1-109 may contain one or more chiral centres, and exist in different optically active forms. When compounds of formulas 1-109 contain one chiral centre, the compounds exist in two enantiomeric forms and the present invention includes both enantiomers and mixtures of enantiomers, such as racemic mixtures. The enantiomers may be resolved by methods known to those skilled in the art, for example by formation of diastereoisomeric salts which may be separated, for example, by crystallization; formation of diastereoisomeric derivatives or complexes which may be separated, for example, by crystallization, gas-liquid or liquid chromatography; selective reaction of one enantiomer with an enantiomer-specific reagent, for example enzymatic esterification; or gas-liquid or liquid chromatography in a chiral environment, for example on a chiral support for example silica with a bound chiral ligand or in the presence of a chiral solvent. It

-53-

will be appreciated that where the desired enantiomer is converted into another chemical entity by one of the separation procedures described above, a further step is required to liberate the desired enantiomeric form. Alternatively, specific enantiomers may be synthesized by asymmetric synthesis using optically active reagents, substrates, catalysts or solvents, or by converting one enantiomer into the other by asymmetric transformation.

When a compound of formulas 1-109 contains more than one chiral centre it may exist in diastereoisomeric forms. The diastereoisomeric pairs may be separated by methods known to those skilled in the art, for example chromatography or crystallization and the individual enantiomers within each pair may be separated as described above. The present invention includes each diastereoisomer of compounds of formulas 1-109 and mixtures thereof.

Certain compounds of formulas 1-109 may exist in different tautomeric forms or as different geometric isomers, and the present invention includes each tautomer and/or geometric isomer of compounds of formulas 1-109 and mixtures thereof.

Certain compounds of formulas 1-109 may exist in different stable conformational forms which may be separable. Torsional asymmetry due to restricted rotation about an asymmetric single bond, for example because of steric hindrance or ring strain, may permit separation of different conformers. The present invention includes each conformational isomer of compounds of formulas 1-109 and mixtures thereof.

Certain compounds of formulas 1-109 may exist in zwitterionic form and the present invention includes each zwitterionic form of compounds of formulas 1-109 and mixtures thereof. Heteroaromatic groups, as used herein, include heteroaryl ring systems (e.g., for purposes of exemplification, which should not be construed as limiting the scope of this invention: thienyl, pyridyl, pyrazole, isoxazolyl, thiadiazolyl, oxadiazolyl, indazolyl, furans, pyrroles, imidazoles, pyrazoles, triazoles, pyrimidines, pyrazines, thiazoles, isothiazoles, oxazolyl or tetrazoles) and heteroaryl ring systems in which a carbocyclic aromatic ring, carbocyclic non-aromatic ring or heteroaryl ring is fused to one or more other heteroaryl rings (e.g.,

for purposes of exemplification, which should not be construed as limiting the scope of this invention: benzo(b)thienyl, benzimidazolyl, benzoxazolyl, benzothiazolyl, benzothiadiazolyl, benzoxadiazolyl, indole, tetrahydroindole, azaindole, indazole, quinoline, imidazopyridine, quinazoline purine, pyrrolo[2,3-d]pyrimidine, pyrazolo[3,4-d]pyrimidine) and their N-oxides. Substituted heteroaryl groups are preferably substituted with one or more substituents each independently selected from the group consisting of a halogen, hydroxy, alkyl, alkoxy, alkyl-O-C(O)-, alkoxyalkyl, a heterocycloalkyl group, optionally substituted phenyl, nitro, amino, mono-substituted amino or di-substituted amino.

10 A heterocyclic (heterocyclyl) group, as used herein, refers to both heteroaryl groups and heterocycloalkyl groups.

 A heterobicyclic group, as used herein, refers to a bicyclic group having one or more heteroatoms, which is saturated, partially unsaturated or unsaturated.

 An arylalkyl group, as used herein, is an aromatic substituent that is linked to a compound by an aliphatic group having from one to about six carbon atoms. A preferred arylalkyl group is a benzyl group

 An heteroaralkyl group, as used herein, is a heteroaromatic substituent that is linked to a compound by an aliphatic group having from one to about six carbon atoms.

20 A heterocycloalkyl group, as used herein, is a non-aromatic ring system that has 3 to 8 atoms and includes at least one heteroatom, such as nitrogen, oxygen, or sulfur. As used herein, aliphatic groups or notations such as "(C₀-C₆)" include straight chained, branched or cyclic hydrocarbons which are completely saturated or which contain one or more units of unsaturation. When the group is a C₀ it means that the moiety is not present or in other words is a bond.

 As used herein, aromatic groups (or aryl groups) include aromatic carbocyclic ring systems (e.g. phenyl) and fused polycyclic aromatic ring systems (e.g. naphthyl and 1,2,3,4-tetrahydronaphthyl).

 As used herein, the term "natural amino acid" refers to the twenty-three natural amino acids known in the art, which are as follows (denoted by their three letter acronym): Ala, Arg, Asn, Asp, Cys, Cys-Cys, Glu, Gln, Gly, His, Hyl, Hyp,

-55-

Ile, Leu, Lys, Met, Phe, Pro, Ser, Thr, Trp, Tyr, and Val. The term non-natural amino acid refers to compounds of the formula $\text{NH}_2\text{-(C(X))}_n\text{-COOH}$, which are alpha- (when n is 1) or beta- (when n is 2) amino acids where X for each occurrence is independently any side chain moiety recognized by those skilled in the art;

5 examples of non-natural amino acids include, but are not limited to: hydroxyproline, homoproline, 4-amino-phenylalanine, β -(2-naphthyl)alanine, norleucine, cyclohexylalanine, β -(3-pyridinyl)alanine, β -(4-pyridinyl)alanine, α -aminoisobutyric acid, urocanic acid, N,N-tetramethylamidino-histidine, N-methyl-alanine, N-methyl-glycine, N-methyl-glutamic acid, *tert*-butylglycine, α -

10 aminobutyric acid, *tert*-butylalanine, ornithine, α -aminoisobutyric acid, β -alanine, γ -aminobutyric acid, 5-aminovaleric acid, 12-aminododecanoic acid, 2-aminoindanc-2-carboxylic acid, etc. and the derivatives thereof, especially where the amine nitrogen has been mono- or di-alkylated.

As used herein, many moieties or substituents are termed as being either

15 "substituted or unsubstituted" or "optionally substituted". When a moiety is modified by one of these terms, it denotes that any portion of the moiety that is known to one skilled in the art as being available for substitution can be substituted, which includes one or more substituents, where if more than one substituent then each substituent is independently selected. Such means for substitution are well-

20 known in the art and/or taught by the instant disclosure. For purposes of exemplification, which should not be construed as limiting the scope of this invention, some examples of groups that are substituents are: alkyl groups (which itself can also be substituted, such as CF_3), alkoxy group (which itself can be substituted, such as OCF_3), a halogen or halo group (F, Cl, Br, I), hydroxy, nitro,

25 oxo, CN, COH, COOH, amino, N-alkylamino or N,N-dialkylamino (in which the alkyl groups can also be substituted), esters ($-\text{C}(\text{O})-\text{OR}$, where R is groups such as alkyl, aryl, etc., which can be substituted), aryl (most preferred is phenyl, which can be substituted) and arylalkyl (which can be substituted).

The compounds of this invention have antiangiogenic properties. These

30 antiangiogenic properties are due at least in part to the inhibition of protein tyrosine kinases essential for angiogenic processes. For this reason, these compounds can be

-56-

used as active agents against such disease states as arthritis, atherosclerosis, restenosis, psoriasis, hemangiomas, myocardial angiogenesis, coronary and cerebral collaterals, ischemic limb angiogenesis, ischemia/reperfusion injury, wound healing, peptic ulcer Helicobacter related diseases, virally-induced angiogenic disorders, fractures, Crow-Fukase syndrome (POEMS), preeclampsia, menometrorrhagia, cat scratch fever, rubeosis, neovascular glaucoma and retinopathies such as those associated with diabetic retinopathy, retinopathy of prematurity, or age-related macular degeneration. In addition, some of these compounds can be used as active agents against solid tumors, malignant ascites, von Hippel Lindau disease, hematopoietic cancers and hyperproliferative disorders such as thyroid hyperplasia (especially Grave's disease), and cysts (such as hypervascularity of ovarian stroma characteristic of polycystic ovarian syndrome (Stein-Leventhal syndrome) and polycystic kidney disease since such diseases require a proliferation of blood vessel cells for growth and/or metastasis.

Further, some of these compounds can be used as active agents against burns, chronic lung disease, stroke, polyps, anaphylaxis, chronic and allergic inflammation, delayed-type hypersensitivity, ovarian hyperstimulation syndrome, brain tumor-associated cerebral edema, high-altitude, trauma or hypoxia induced cerebral or pulmonary edema, ocular and macular edema, ascites, glomerulonephritis and other diseases where vascular hyperpermeability, effusions, exudates, protein extravasation, or edema is a manifestation of the disease. The compounds will also be useful in treating disorders in which protein extravasation leads to the deposition of fibrin and extracellular matrix, promoting stromal proliferation (e.g. keloid, fibrosis, cirrhosis and carpal tunnel syndrome). Increased VEGF production potentiates inflammatory processes such as monocyte recruitment and activation. The compounds of this invention will also be useful in treating inflammatory disorders such as inflammatory bowel disease (IBD) and Crohn's disease.

VEGF's are unique in that they are the only angiogenic growth factors known to contribute to vascular hyperpermeability and the formation of edema. Indeed, vascular hyperpermeability and edema that is associated with the expression or administration of many other growth factors appears to be mediated via VEGF

production. Inflammatory cytokines stimulate VEGF production. Hypoxia results in a marked upregulation of VEGF in numerous tissues, hence situations involving infarct, occlusion, ischemia, anemia, or circulatory impairment typically invoke VEGF/VPF mediated responses. Vascular hyperpermeability, associated edema, altered transendothelial exchange and macromolecular extravasation, which is often accompanied by diapedesis, can result in excessive matrix deposition, aberrant stromal proliferation, fibrosis, etc. Hence, VEGF-mediated hyperpermeability can significantly contribute to disorders with these etiologic features.

Because blastocyst implantation, placental development and embryogenesis are angiogenesis dependent, certain compounds of the invention are useful as contraceptive agents and antifertility agents.

It is envisaged that the disorders listed above are mediated to a significant extent by protein tyrosine kinase activity involving the KDR/VEGFR-2 and/or the Flt-1/VEGFR-1 and/or TIE-2 tyrosine kinases. By inhibiting the activity of these tyrosine kinases, the progression of the listed disorders is inhibited because the angiogenic or vascular hyperpermeability component of the disease state is severely curtailed. The action of certain compounds of this invention, by their selectivity for specific tyrosine kinases, result in a minimization of side effects that would occur if less selective tyrosine kinase inhibitors were used. Certain compounds of the invention are also effective inhibitors of FGFR, PDGFR, c-Met and IGF-1-R. These receptor kinases can directly or indirectly potentiate angiogenic and hyperproliferative responses in various disorders, hence their inhibition can impede disease progression.

Tie-2 (TEK) is a member of a recently discovered family of endothelial cell specific receptor tyrosine kinases which is involved in critical angiogenic processes, such as vessel branching, sprouting, remodeling, maturation and stability. Tie-2 is the first mammalian receptor tyrosine kinase for which both agonist ligand(s) (e.g., Angiopoietin1 ("Ang1")), which stimulates receptor autophosphorylation and signal transduction), and antagonist ligand(s) (e.g., Angiopoietin2 ("Ang2")), have been identified. Knock-out and transgenic manipulation of the expression of Tie-2 and its ligands indicates tight spatial and temporal control of Tie-2 signaling is essential for

-58-

the proper development of new vasculature. The current model suggests that stimulation of Tie-2 kinase by the Ang1 ligand is directly involved in the branching, sprouting and outgrowth of new vessels, and recruitment and interaction of periendothelial support cells important in maintaining vessel integrity and inducing quiescence. The absence of Ang1 stimulation of Tie-2 or the inhibition of Tie-2 autophosphorylation by Ang2, which is produced at high levels at sites of vascular regression, may cause a loss in vascular structure and matrix contacts resulting in endothelial cell death, especially in the absence of growth/survival stimuli. The situation is however more complex, since at least two additional Tie-2 ligands (Ang3 and Ang4) have recently been reported, and the capacity for heterooligomerization of the various agonistic and antagonistic angiopoietins, thereby modifying their activity, has been demonstrated. Targeting Tie-2 ligand-receptor interactions as an antiangiogenic therapeutic approach is thus less favored and a kinase inhibitory strategy preferred.

The soluble extracellular domain of Tie-2 ("ExTek") can act to disrupt the establishment of tumor vasculature in a breast tumor xenograft and lung metastasis models and in tumor-cell mediated ocular neovascularization. By adenoviral infection, the *in vivo* production of mg/ml levels ExTek in rodents may be achieved for 7-10 days with no adverse side effects. These results suggest that disruption of Tie-2 signaling pathways in normal healthy animals may be well tolerated. These Tie-2 inhibitory responses to ExTek may be a consequence sequestration of ligand(s) and/or generation of a nonproductive heterodimer with full-length Tie-2.

Recently, significant upregulation of Tie-2 expression has been found within the vascular synovial pannus of arthritic joints of humans, consistent with a role in the inappropriate neovascularization. This finding suggests that Tie-2 plays a role in the progression of rheumatoid arthritis. Point mutations producing constitutively activated forms of Tie-2 have been identified in association with human venous malformation disorders. Tie-2 inhibitors are, therefore, useful in treating such disorders, and in other situations of inappropriate neovascularization.

The compounds of this invention have inhibitory activity against protein kinases. That is, these compounds modulate signal transduction by protein kinases.

-59-

Compounds of this invention inhibit protein kinases from serine/threonine and tyrosine kinase classes. In particular, these compounds selectively inhibit the activity of the KDR/FLK-1/VEGFR-2 tyrosine kinases. Certain compounds of this invention also inhibit the activity of additional tyrosine kinases such as Flt-1/VEGFR-1, Flt-4/VEGFR-3, Tie-1, Tie-2, FGFR, PDGFR, IGF-1R, c-Met, Src-subfamily kinases such as Lck, hck, fgr, Src, fyn, yes, etc. Additionally, some compounds of this invention significantly inhibit serine/threonine kinases such as PKC, MAP kinases, erk, CDKs, Plk-1, or Raf-1 which play an essential role in cell proliferation and cell-cycle progression. The potency and specificity of the generic compounds of this invention towards a particular protein kinase can often be altered and optimized by variations in the nature, number and arrangement of the substituents (i.e., R₁, R₂, R₃, A and ring 1) and conformational restrictions. In addition the metabolites of certain compounds may also possess significant protein kinase inhibitory activity.

The compounds of this invention, when administered to individuals in need of such compounds, inhibit vascular hyperpermeability and the formation of edema in these individuals. These compounds act, it is believed, by inhibiting the activity of KDR tyrosine kinase which is involved in the process of vascular hyperpermeability and edema formation. The KDR tyrosine kinase may also be referred to as FLK-1 tyrosine kinase, NYK tyrosine kinase or VEGFR-2 tyrosine kinase. KDR tyrosine kinase is activated when vascular endothelial cell growth factor (VEGF) or another activating ligand (such as VEGF-C, VEGF-D, VEGF-E or HIV Tat protein) binds to a KDR tyrosine kinase receptor which lies on the surface of vascular endothelial cells. Following such KDR tyrosine kinase activation, hyperpermeability of the blood vessels occurs and fluid moves from the blood stream past the blood vessel walls into the interstitial spaces, thereby forming an area of edema. Diapedesis also often accompanies this response. Similarly, excessive vascular hyperpermeability can disrupt normal molecular exchange across the endothelium in critical tissues and organs (e.g., lung and kidney), thereby causing macromolecular extravasation and deposition. Following this acute response to KDR stimulation which is believed to facilitate the subsequent

-60-

angiogenic process, prolonged KDR tyrosine kinase stimulation results in the proliferation and chemotaxis of vascular endothelial cells and formation of new vessels. By inhibiting KDR tyrosine kinase activity, either by blocking the production of the activating ligand, by blocking the activating ligand binding to the KDR tyrosine kinase receptor, by preventing receptor dimerization and transphosphorylation, by inhibiting the enzyme activity of the KDR tyrosine kinase (inhibiting the phosphorylation function of the enzyme) or by some other mechanism that interrupts its downstream signaling (D. Mukhopedhyay *et al.*, *Cancer Res.* 58:1278-1284 (1998) and references therein), hyperpermeability, as well as associated extravasation, subsequent edema formation and matrix deposition, and angiogenic responses, may be inhibited and minimized.

One group of preferred compounds of this invention have the property of inhibiting KDR tyrosine kinase activity without significantly inhibiting Flt-1 tyrosine kinase activity (Flt-1 tyrosine kinase is also referred to as VEGFR-1 tyrosine kinase). Both KDR tyrosine kinase and Flt-1 tyrosine kinase are activated by VEGF binding to KDR tyrosine kinase receptors and to Flt-1 tyrosine kinase receptors, respectively. Certain preferred compounds of this invention are unique because they inhibit the activity of one VEGF-receptor tyrosine kinase (KDR) that is activated by activating ligands but do not inhibit other receptor tyrosine kinases, such as Flt-1, that are also activated by certain activating ligands. In this manner, certain preferred compounds of this invention are, therefore, selective in their tyrosine kinase inhibitory activity.

In one embodiment, the present invention provides a method of treating a protein kinase-mediated condition in a patient, comprising administering to the patient a therapeutically or prophylactically effective amount of one or more compounds of Formulas 1-109.

A "protein kinase-mediated condition" or a "condition mediated by protein kinase activity" is a medical condition, such as a disease or other undesirable physical condition, the genesis or progression of which depends, at least in part, on the activity of at least one protein kinase. The protein kinase can be, for example, a protein tyrosine kinase or a protein serine/threonine kinase.

-61-

The patient to be treated can be any animal, and is preferably a mammal, such as a domesticated animal or a livestock animal. More preferably, the patient is a human.

A "therapeutically effective amount" is an amount of a compound of
5 Formulas 1-109 or a combination of two or more such compounds, which inhibits, totally or partially, the progression of the condition or alleviates, at least partially, one or more symptoms of the condition. A therapeutically effective amount can also be an amount which is prophylactically effective. The amount which is
10 therapeutically effective will depend upon the patient's size and gender, the condition to be treated, the severity of the condition and the result sought. For a given patient, a therapeutically effective amount can be determined by methods known to those of skill in the art.

The method of the present invention is useful in the treatment of protein kinase-mediated conditions, such as any of the conditions described above. In one
15 embodiment, the protein kinase-mediated condition is characterized by undesired angiogenesis, edema, or stromal deposition. For example, the condition can be one or more more ulcers, such as ulcers caused by bacterial or fungal infections, Mooren ulcers and ulcerative colitis. The condition can also be due to a microbial infection, such as Lyme disease, sepsis, septic shock or infections by Herpes simplex, Herpes
20 Zoster, human immunodeficiency virus, protozoa, toxoplasmosis or parapoxvirus; an angiogenic disorders, such as von Hippel Lindau disease, polycystic kidney disease, pemphigoid, Paget's disease and psoriasis; a reproductive condition, such as endometriosis, ovarian hyperstimulation syndrome, preeclampsia or menometrorrhagia; a fibrotic and edemic condition, such as sarcoidosis, fibrosis,
25 cirrhosis, thyroiditis, hyperviscosity syndrome systemic, Osler-Weber-Rendu disease, chronic occlusive pulmonary disease, asthma, and edema following burns, trauma, radiation, stroke, hypoxia or ischemia; or an inflammatory/immunologic condition, such as systemic lupus, chronic inflammation, glomerulonephritis, synovitis, inflammatory bowel disease, Crohn's disease, rheumatoid arthritis,
30 osteoarthritis, multiple sclerosis and graft rejection. Suitable protein kinase-mediated conditions also include sickle cell anaemia, osteoporosis, osteopetrosis,

-62-

tumor-induced hypercalcemia and bone metastases. Additional protein kinase-mediated conditions which can be treated by the method of the present invention include ocular conditions such as ocular and macular edema, ocular neovascular disease, scleritis, radial keratotomy, uveitis, vitritis, myopia, optic pits, chronic
5 retinal detachment, post-laser complications, conjunctivitis, Stargardt's disease and Eales disease, in addition to retinopathy and macular degeneration.

The compounds of the present invention are also useful in the treatment of cardiovascular conditions such as atherosclerosis, restenosis, vascular occlusion and carotid obstructive disease.

10 The compounds of the present invention are also useful in the treatment of cancer related indications such as solid tumors, sarcomas (especially Ewing's sarcoma and osteosarcoma), retinoblastoma, rhabdomyosarcomas, neuroblastoma, hematopoietic malignancies, including leukaemia and lymphoma, tumor-induced pleural or pericardial effusions, and malignant ascites.

15 The compounds of the present invention are also useful in the treatment of Crow-Fukase (POEMS) syndrome and diabetic conditions such as glaucoma, diabetic retinopathy and microangiopathy.

The Src, Tec, Jak, Map, Csk, NF κ B and Syk families of kinases play pivotal roles in the regulation of immune function. The Src family currently includes Fyn,
20 Lck, Fgr, Fes, Lyn, Src, Yrk, Fyk, Yes, Hck, and Blk. The Syk family is currently understood to include only Zap and Syk. The TEC family includes Tec, Btk, Rlk and Itk. The Janus family of kinases is involved in the transduction of growth factor and proinflammatory cytokine signals through a number of receptors. Although BTK and ITK, members of the Tec family of kinases, play a less well understood role in
25 immunobiology, their modulation by an inhibitor may prove therapeutically beneficial. The Csk family is currently understood to include Csk and Chk. The kinases RIP, IRAK-1, IRAK-2, NIK, p38 MAP kinases, Jnk, IKK-1 and IKK-2 are involved in the signal transduction pathways for key pro-inflammatory cytokines, such as TNF and IL-1. By virtue of their ability to inhibit one or more of these
30 kinases, compounds of formulas 1-109 may function as immunomodulatory agents useful for the maintenance of allografts, the treatment of autoimmune disorders and

treatment of sepsis and septic shock. Through their ability to regulate the migration or activation of T cells, B-cells, mast cells, monocytes and neutrophils, these compounds could be used to treat such autoimmune diseases and sepsis. Prevention of transplant rejection, either host versus graft for solid organs or graft versus host
5 for bone marrow, are limited by the toxicity of currently available immunosuppressive agents and would benefit from an efficacious drug with improved therapeutic index. Gene targeting experiments have demonstrated the essential role of Src in the biology of osteoclasts, the cells responsible for bone resorption. Compounds of formulas 1-109, through their ability to regulate Src, may
10 also be useful in the treatment of osteoporosis, osteopetrosis, Paget's disease, tumor-induced hypercalcemia and in the treatment of bone metastases.

A number of protein kinases have been demonstrated to be protooncogenes. Chromosome breakage (at the ltk kinase break point on chromosome 5), translocation as in the case of the Abl gene with BCR (Philadelphia chromosome),
15 truncation in instances such as c-Kit or EGFR, or mutation (e.g., Met) result in the creation of dysregulated proteins converting them from protooncogene to oncogene products. In other tumors, oncogenesis is driven by an autocrine or paracrine ligand/growth factor receptor interactions. Members of the src-family kinases are typically involved in downstream signal transduction thereby potentiating the
20 oncogenesis and themselves may become oncogenic by over-expression or mutation. By inhibiting the protein kinase activity of these proteins the disease process may be disrupted. Vascular restenosis may involve FGF and/or PDGF - promoted smooth muscle and endothelial cell proliferation. The ligand stimulation of FGFR, PDGFR, IGF1-R and c-Met *in vivo* is proangiogenic, and potentiates angiogenesis dependent
25 disorders. Inhibition of FGFR, PDGFR, c-Met, or IGF1-R kinase activities individually or in combination may be an efficacious strategy for inhibiting these phenomena. Thus compounds of formulas 1-109 which inhibit the kinase activity of normal or aberrant c-kit, c-met, c-fms, src-family members, EGFR, erbB2, erbB4, BCR-Abl, PDGFR, FGFR, IGF1-R and other receptor or cytosolic tyrosine kinases
30 may be of value in the treatment of benign and neoplastic proliferative diseases.

-64-

In many pathological conditions (for example, solid primary tumors and metastases, Kaposi's sarcoma, rheumatoid arthritis, blindness due to inappropriate ocular neovascularization, psoriasis and atherosclerosis) disease progression is contingent upon persistent angiogenesis. Polypeptide growth factors often produced by the disease tissue or associated inflammatory cells, and their corresponding endothelial cell specific receptor tyrosine kinases (e.g., KDR/VEGFR-2, Flt-1/VEGFR-1, Tie-2/Tek and Tie) are essential for the stimulation of endothelial cell growth, migration, organization, differentiation and the establishment of the requisite new functional vasculature. As a result of the vascular permeability factor activity of VEGF in mediating vascular hyperpermeability, VEGF-stimulation of a VEGFR kinase is also believed to play an important role in the formation of tumor ascites, cerebral and pulmonary edema, pleural and pericardial effusions, delayed-type hypersensitivity reactions, tissue edema and organ dysfunction following trauma, burns, ischemia, diabetic complications, endometriosis, adult respiratory distress syndrome (ARDS), post-cardiopulmonary bypass-related hypotension and hyperpermeability, and ocular edema leading to glaucoma or blindness due to inappropriate neovascularization. In addition to VEGF, recently identified VEGF-C and VEGF-D, and virally-encoded VEGF-E or HIV-Tat protein can also cause a vascular hyperpermeability response through the stimulation of a VEGFR kinase. KDR/VEGFR-2 and/or Tie-2 are expressed also in a select population of hematopoietic stem cells. Certain members of this population are pluripotent in nature and can be stimulated with growth factors to differentiate into endothelial cells and participate in vasculogenetic angiogenic processes. For this reason these have been called Endothelial Progenitor Cells (EPCs) (*J. Clin. Investig.* 103 : 1231-1236 (1999)). In some progenitors, Tie-2 may play a role in their recruitment, adhesion, regulation and differentiation (*Blood* , 4317-4326 (1997)). Certain agents according to formulas 1-109 capable of blocking the kinase activity of endothelial cell specific kinases could therefore inhibit disease progression involving these situations.

Vascular destabilization of the antagonist ligand of Tie-2 (Ang2) is believed to induce an unstable "plastic" state in the endothelium. In the presence of high

VEGF levels a robust angiogenic response may result; however, in the absence of VEGF or a VEGF-related stimulus, frank vessel regression and endothelial apoptosis can occur (Genes and Devel. 13: 1055-1066 (1999)). In an analogous manner a Tie-2 kinase inhibitor can be proangiogenic or antiangiogenic in the presence or absence
5 of a VEGF-related stimulus, respectively.

The compounds of formulas 1-109 or a salt thereof or pharmaceutical compositions containing a therapeutically effective amount thereof may be used in the treatment of protein kinase-mediated conditions, such as benign and neoplastic proliferative diseases and disorders of the immune system, as described above. For
10 example, such diseases include autoimmune diseases, such as rheumatoid arthritis, thyroiditis, type 1 diabetes, multiple sclerosis, sarcoidosis, inflammatory bowel disease, Crohn's disease, myasthenia gravis and systemic lupus erythematosus; psoriasis, organ transplant rejection (eg. kidney rejection, graft versus host disease), benign and neoplastic proliferative diseases, human cancers such as lung, breast,
15 stomach, bladder, colon, pancreas, ovarian, prostate and rectal cancer and hematopoietic malignancies (leukemia and lymphoma), and diseases involving inappropriate vascularization for example diabetic retinopathy, retinopathy of prematurity, choroidal neovascularization due to age-related macular degeneration, and infantile hemangiomas in human beings. In addition, such inhibitors may be
20 useful in the treatment of disorders involving VEGF mediated edema, ascites, effusions, and exudates, including for example macular edema, cerebral edema, acute lung injury and adult respiratory distress syndrome (ARDS).

The compounds of the present invention may also be useful in the prophylaxis of the above diseases.

25 It is envisaged that the disorders listed above are mediated to a significant extent by protein tyrosine kinase activity involving the VEGF receptors (e.g. KDR, Flt-1 and/or Tie-2). By inhibiting the activity of these receptor tyrosine kinases, the progression of the listed disorders is inhibited because the angiogenic component of the disease state is severely curtailed. The action of the compounds of this
30 invention, by their selectivity for specific tyrosine kinases, result in a minimization of side effects that would occur if less selective tyrosine kinase inhibitors were used.

-66-

In another aspect the present invention provides compounds of formulas 1-109 as defined initially above for use as medicaments, particularly as inhibitors of protein kinase activity for example tyrosine kinase activity, serine kinase activity and threonine kinase activity. In yet another aspect the present invention provides the use of compounds of formulas 1-109 as defined initially above in the manufacture of a medicament for use in the inhibition of protein kinase activity.

In this invention, the following definitions are applicable:

"Physiologically acceptable salts" refers to those salts which retain the biological effectiveness and properties of the free bases and which are obtained by reaction with inorganic acids such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid or organic acids such as sulfonic acid, carboxylic acid, organic phosphoric acid, methanesulfonic acid, ethanesulfonic acid, p-toluenesulfonic acid, salicylic acid, lactic acid, tartaric acid and the like.

15 Pharmaceutical Formulations

The compounds of this invention can be administered to a human patient by themselves or in pharmaceutical compositions where they are mixed with suitable carriers or excipient(s) at doses to treat or ameliorate vascular hyperpermeability, edema and associated disorders. Mixtures of these compounds can also be administered to the patient as a simple mixture or in suitable formulated pharmaceutical compositions. A therapeutically effective dose further refers to that amount of the compound or compounds sufficient to result in the prevention or attenuation of inappropriate neovascularization, progression of hyperproliferative disorders, edema, VEGF-associated hyperpermeability and/or VEGF-related hypotension. Techniques for formulation and administration of the compounds of the instant application may be found in "Remington's Pharmaceutical Sciences," Mack Publishing Co., Easton, PA, latest edition.

Routes of Administration

Suitable routes of administration may, for example, include oral, eyedrop, rectal, transmucosal, topical, or intestinal administration; parenteral delivery,

-67-

including intramuscular, subcutaneous, intramedullary injections, as well as intrathecal, direct intraventricular, intravenous, intraperitoneal, intranasal, or intraocular injections.

Alternatively, one may administer the compound in a local rather than a systemic manner, for example, via injection of the compound directly into an edematous site, often in a depot or sustained release formulation.

Furthermore, one may administer the drug in a targeted drug delivery system, for example, in a liposome coated with endothelial cell-specific antibody.

10 Composition/Formulation

The pharmaceutical compositions of the present invention may be manufactured in a manner that is itself known, e.g., by means of conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping or lyophilizing processes.

15 Pharmaceutical compositions for use in accordance with the present invention thus may be formulated in conventional manner using one or more physiologically acceptable carriers comprising excipients and auxiliaries which facilitate processing of the active compounds into preparations which can be used pharmaceutically. Proper formulation is dependent upon the route of administration
20 chosen.

For injection, the agents of the invention may be formulated in aqueous solutions, preferably in physiologically compatible buffers such as Hanks's solution, Ringer's solution, or physiological saline buffer. For transmucosal administration, penetrants appropriate to the barrier to be permeated are used in the formulation.

25 Such penetrants are generally known in the art.

For oral administration, the compounds can be formulated readily by combining the active compounds with pharmaceutically acceptable carriers well known in the art. Such carriers enable the compounds of the invention to be formulated as tablets, pills, dragees, capsules, liquids, gels, syrups, slurries, suspensions and the like, for oral ingestion by a patient to be treated.

30 Pharmaceutical preparations for oral use can be obtained by combining the active

-68-

compound with a solid excipient, optionally grinding a resulting mixture, and processing the mixture of granules, after adding suitable auxiliaries, if desired, to obtain tablets or dragee cores. Suitable excipients are, in particular, fillers such as sugars, including lactose, sucrose, mannitol, or sorbitol; cellulose preparations such as, for example, maize starch, wheat starch, rice starch, potato starch, gelatin, gum tragacanth, methyl cellulose, hydroxypropylmethyl-cellulose, sodium carboxymethylcellulose, and/or polyvinylpyrrolidone (PVP). If desired, disintegrating agents may be added, such as the cross-linked polyvinyl pyrrolidone, agar, or alginic acid or a salt thereof such as sodium alginate.

- 10 Dragee cores are provided with suitable coatings. For this purpose, concentrated sugar solutions may be used, which may optionally contain gum arabic, talc, polyvinyl pyrrolidone, carbopol gel, polyethylene glycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. Dyestuffs or pigments may be added to the tablets or dragee coatings for
- 15 identification or to characterize different combinations of active compound doses.

- Pharmaceutical preparations which can be used orally include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a plasticizer, such as glycerol or sorbitol. The push-fit capsules can contain the active ingredients in admixture with filler such as lactose, binders such as starches, and/or
- 20 lubricants such as talc or magnesium stearate and, optionally, stabilizers. In soft capsules, the active compounds may be dissolved or suspended in suitable liquids, such as fatty oils, liquid paraffin, or liquid polyethylene glycols. In addition, stabilizers may be added. All formulations for oral administration should be in dosages suitable for such administration.

- 25 For buccal administration, the compositions may take the form of tablets or lozenges formulated in conventional manner.

- For administration by inhalation, the compounds for use according to the present invention are conveniently delivered in the form of an aerosol spray presentation from pressurized packs or a nebuliser, with the use of a suitable
- 30 propellant, e.g., dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide or other suitable gas. In the case of

pressurized aerosol the dosage unit may be determined by providing a valve to deliver a metered amount. Capsules and cartridges of e.g. gelatin for use in an inhaler or insufflator may be formulated containing a powder mix of the compound and a suitable powder base such as lactose or starch.

5 The compounds can be formulated for parenteral administration by injection, e.g. bolus injection or continuous infusion. Formulations for injection may be presented in unit dosage form, e.g. in ampoules or in multi-dose containers, with an added preservative. The compositions may take such forms as suspensions, solutions or emulsions in oily or aqueous vehicles, and may contain formulatory
10 agents such as suspending, stabilizing and/or dispersing agents.

Pharmaceutical formulations for parenteral administration include aqueous solutions of the active compounds in water-soluble form. Additionally, suspensions of the active compounds may be prepared as appropriate oily injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils such as sesame oil, or
15 synthetic fatty acid esters, such as ethyl oleate or triglycerides, or liposomes. Aqueous injection suspensions may contain substances which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran. Optionally, the suspension may also contain suitable stabilizers or agents which increase the solubility of the compounds to allow for the preparation of highly
20 concentrated solutions.

Alternatively, the active ingredient may be in powder form for constitution with a suitable vehicle, e.g., sterile pyrogen-free water, before use.

The compounds may also be formulated in rectal compositions such as suppositories or retention enemas, e.g., containing conventional suppository bases
25 such as cocoa butter or other glycerides.

In addition to the formulations described previously, the compounds may also be formulated as a depot preparation. Such long acting formulations may be administered by implantation (for example subcutaneously or intramuscularly or by intramuscular injection). Thus, for example, the compounds may be formulated with
30 suitable polymeric or hydrophobic materials (for example as an emulsion in an acceptable oil) or ion exchange resins, or as sparingly soluble derivatives, for

-70-

example, as a sparingly soluble salt.

An example of a pharmaceutical carrier for the hydrophobic compounds of the invention is a cosolvent system comprising benzyl alcohol, a nonpolar surfactant, a water-miscible organic polymer, and an aqueous phase. The cosolvent system may be the VPD co-solvent system. VPD is a solution of 3% w/v benzyl alcohol, 8% w/v of the nonpolar surfactant polysorbate 80, and 65% w/v polyethylene glycol 300, made up to volume in absolute ethanol. The VPD co-solvent system (VPD:5W) consists of VPD diluted 1:1 with a 5% dextrose in water solution. This co-solvent system dissolves hydrophobic compounds well, and itself produces low toxicity upon systemic administration. Naturally, the proportions of a co-solvent system may be varied considerably without destroying its solubility and toxicity characteristics. Furthermore, the identity of the co-solvent components may be varied: for example, other low-toxicity nonpolar surfactants may be used instead of polysorbate 80; the fraction size of polyethylene glycol may be varied; other biocompatible polymers may replace polyethylene glycol, e.g. polyvinyl pyrrolidone; and other sugars or polysaccharides may substitute for dextrose.

Alternatively, other delivery systems for hydrophobic pharmaceutical compounds may be employed. Liposomes and emulsions are well known examples of delivery vehicles or carriers for hydrophobic drugs. Certain organic solvents such as dimethylsulfoxide also may be employed, although usually at the cost of greater toxicity. Additionally, the compounds may be delivered using a sustained-release system, such as semipermeable matrices of solid hydrophobic polymers containing the therapeutic agent. Various sustained-release materials have been established and are well known by those skilled in the art. Sustained-release capsules may, depending on their chemical nature, release the compounds for a few weeks up to over 100 days. Depending on the chemical nature and the biological stability of the therapeutic reagent, additional strategies for protein stabilization may be employed.

The pharmaceutical compositions also may comprise suitable solid or gel phase carriers or excipients. Examples of such carriers or excipients include but are not limited to calcium carbonate, calcium phosphate, various sugars, starches,

-71-

cellulose derivatives, gelatin, and polymers such as polyethylene glycols.

Many of the compounds of the invention may be provided as salts with pharmaceutically compatible counterions. Pharmaceutically compatible salts may be formed with many acids, including but not limited to hydrochloric, sulfuric, acetic, lactic, tartaric, malic, succinic, etc. Salts tend to be more soluble in aqueous or other protonic solvents than are the corresponding free base forms.

Effective Dosage

Pharmaceutical compositions suitable for use in the present invention include compositions wherein the active ingredients are contained in an effective amount to achieve its intended purpose. More specifically, a therapeutically effective amount means an amount effective to prevent development of or to alleviate the existing symptoms of the subject being treated. Determination of the effective amounts is well within the capability of those skilled in the art.

For any compound used in the method of the invention, the therapeutically effective dose can be estimated initially from cellular assays. For example, a dose can be formulated in cellular and animal models to achieve a circulating concentration range that includes the IC_{50} as determined in cellular assays (i.e., the concentration of the test compound which achieves a half-maximal inhibition of a given protein kinase activity). In some cases it is appropriate to determine the IC_{50} in the presence of 3 to 5% serum albumin since such a determination approximates the binding effects of plasma protein on the compound. Such information can be used to more accurately determine useful doses in humans. Further, the most preferred compounds for systemic administration effectively inhibit protein kinase signaling in intact cells at levels that are safely achievable in plasma.

A therapeutically effective dose refers to that amount of the compound that results in amelioration of symptoms in a patient. Toxicity and therapeutic efficacy of such compounds can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, e.g., for determining the maximum tolerated dose (MTD) and the ED_{50} (effective dose for 50% maximal response). The dose ratio between toxic and therapeutic effects is the therapeutic index and it can be expressed

as the ratio between MTD and ED₅₀. Compounds which exhibit high therapeutic indices are preferred. The data obtained from these cell culture assays and animal studies can be used in formulating a range of dosage for use in humans. The dosage of such compounds lies preferably within a range of circulating concentrations that include the ED₅₀ with little or no toxicity. The dosage may vary within this range depending upon the dosage form employed and the route of administration utilized. The exact formulation, route of administration and dosage can be chosen by the individual physician in view of the patient's condition. (See e.g. Fingl *et al.*, 1975, in "The Pharmacological Basis of Therapeutics", Ch. 1 p1). In the treatment of crises, the administration of an acute bolus or an infusion approaching the MTD may be required to obtain a rapid response.

Dosage amount and interval may be adjusted individually to provide plasma levels of the active moiety which are sufficient to maintain the kinase modulating effects, or minimal effective concentration (MEC). The MEC will vary for each compound but can be estimated from *in vitro* data; e.g. the concentration necessary to achieve 50-90% inhibition of protein kinase using the assays described herein. Dosages necessary to achieve the MEC will depend on individual characteristics and route of administration. However, HPLC assays or bioassays can be used to determine plasma concentrations.

Dosage intervals can also be determined using the MEC value. Compounds should be administered using a regimen which maintains plasma levels above the MEC for 10-90% of the time, preferably between 30-90% and most preferably between 50-90% until the desired amelioration of symptoms is achieved. In cases of local administration or selective uptake, the effective local concentration of the drug may not be related to plasma concentration.

The amount of composition administered will, of course, be dependent on the subject being treated, on the subject's weight, the severity of the affliction, the manner of administration and the judgment of the prescribing physician.

Packaging

The compositions may, if desired, be presented in a pack or dispenser device which may contain one or more unit dosage forms containing the active ingredient. The pack may for example comprise metal or plastic foil, such as a blister pack. The pack or dispenser device may be accompanied by instructions for administration. Compositions comprising a compound of the invention formulated in a compatible pharmaceutical carrier may also be prepared, placed in an appropriate container, and labeled for treatment of an indicated condition.

In some formulations it may be beneficial to use the compounds of the present invention in the form of particles of very small size, for example as obtained by fluid energy milling.

The use of compounds of the present invention in the manufacture of pharmaceutical compositions is illustrated by the following description. In this description the term "active compound" denotes any compound of the invention but particularly any compound which is the final product of one of the preceding Examples.

a) Capsules

In the preparation of capsules, 10 parts by weight of active compound and 240 parts by weight of lactose can be de-aggregated and blended. The mixture can be filled into hard gelatin capsules, each capsule containing a unit dose or part of a unit dose of active compound.

b) Tablets

Tablets can be prepared from the following ingredients.

Parts by weight

Active compound	10
Lactose	190
Maize starch	22
Polyvinylpyrrolidone	10
Magnesium stearate	3

The active compound, the lactose and some of the starch can be de-aggregated, blended and the resulting mixture can be granulated with a solution of

-74-

the polyvinyl- pyrrolidone in ethanol. The dry granulate can be blended with the magnesium stearate and the rest of the starch. The mixture is then compressed in a tableting machine to give tablets each containing a unit dose or a part of a unit dose of active compound.

5

c) Enteric coated tablets

Tablets can be prepared by the method described in (b) above. The tablets can be enteric coated in a conventional manner using a solution of 20% cellulose acetate phthalate and 3% diethyl phthalate in ethanol:dichloromethane (1:1).

10

d) Suppositories

In the preparation of suppositories, 100 parts by weight of active compound can be incorporated in 1300 parts by weight of triglyceride suppository base and the mixture formed into suppositories each containing a therapeutically effective amount of active ingredient.

15

In the compositions of the present invention the active compound may, if desired, be associated with other compatible pharmacologically active ingredients. For example, the compounds of this invention can be administered in combination with one or more additional pharmaceutical agents that inhibit or prevent the production of VEGF or angiopoietins, attenuate intracellular responses to VEGF or angiopoietins, block intracellular signal transduction, inhibit vascular hyperpermeability, reduce inflammation, or inhibit or prevent the formation of edema or neovascularization. The compounds of the invention can be administered prior to, subsequent to or simultaneously with the additional pharmaceutical agent, whichever course of administration is appropriate. The additional pharmaceutical agents include but are not limited to anti-edemic steroids, NSAIDS, ras inhibitors, anti-TNF agents, anti-IL1 agents, antihistamines, PAF-antagonists, COX-1 inhibitors, COX-2 inhibitors, NO synthase inhibitors, Akt/PTB inhibitors, IGF-1R inhibitors, PKC inhibitors and PI3 kinase inhibitors. The compounds of the invention and the additional pharmaceutical agents act either additively or synergistically. Thus, the administration of such a combination of substances that

20

25

30

-75-

inhibit angiogenesis, vascular hyperpermeability and/or inhibit the formation of edema can provide greater relief from the deletrious effects of a hyperproliferative disorder, angiogenesis, vascular hyperpermeability or edema than the administration of either substance alone. In the treatment of malignant disorders combinations with antiproliferative or cytotoxic chemotherapies or radiation are anticipated.

The present invention also comprises the use of a compound of formulas 1-109 as a medicament.

A further aspect of the present invention provides the use of a compound of formulas 1-109 or a salt thereof in the manufacture of a medicament for treating vascular hyperpermeability, angiogenesis-dependent disorders, proliferative diseases and/or disorders of the immune system in mammals, particularly human beings.

The present invention also provides a method of treating vascular hyperpermeability, inappropriate neovascularization, proliferative diseases and/or disorders of the immune system which comprises the administration of a therapeutically effective amount of a compound of formulas 1-109 to a mammal, particularly a human being, in need thereof.

The *in vitro* potency of compounds in inhibiting these protein kinases may be determined by the procedures detailed below.

The potency of compounds can be determined by the amount of inhibition of the phosphorylation of an exogenous substrate (e.g., synthetic peptide (Z. Songyang *et al.*, *Nature*. 373:536-539) by a test compound relative to control.

KDR Tyrosine Kinase Production Using Baculovirus System:

The coding sequence for the human KDR intra-cellular domain (aa789-1354) was generated through PCR using cDNAs isolated from HUVEC cells. A poly-His6 sequence was introduced at the N-terminus of this protein as well. This fragment was cloned into transfection vector pVL1393 at the Xba 1 and Not 1 site.

Recombinant baculovirus (BV) was generated through co-transfection using the BaculoGold Transfection reagent (PharMingen). Recombinant BV was plaque purified and verified through Western analysis. For protein production, SF-9 cells were grown in SF-900-II medium at 2×10^6 /ml, and were infected at 0.5 plaque

-76-

forming units per cell (MOI). Cells were harvested at 48 hours post infection.

Purification of KDR

SF-9 cells expressing (His)₆KDR(aa789-1354) were lysed by adding 50 ml of
5 Triton X-100 lysis buffer (20 mM Tris, pH 8.0, 137 mM NaCl, 10% glycerol, 1%
Triton X-100, 1mM PMSF, 10μg/ml aprotinin, 1 μg/ml leupeptin) to the cell pellet
from 1L of cell culture. The lysate was centrifuged at 19,000 rpm in a Sorval SS-34
rotor for 30 min at 4°C. The cell lysate was applied to a 5 ml NiCl₂ chelating
sepharose column, equilibrated with 50 mM HEPES, pH7.5, 0.3 M NaCl. KDR was
10 eluted using the same buffer containing 0.25 M imidazole. Column fractions were
analyzed using SDS-PAGE and an ELISA assay (below) which measures kinase
activity. The purified KDR was exchanged into 25mM HEPES, pH7.5, 25mM
NaCl, 5 mM DTT buffer and stored at -80°C.

15 Human Tie-2 Kinase Production and Purification

The coding sequence for the human Tie-2 intra-cellular domain (aa775-1124)
was generated through PCR using cDNAs isolated from human placenta as a
template. A poly-His₆ sequence was introduced at the N-terminus and this construct
was cloned into transfection vector pVL 1939 at the Xba 1 and Not 1 site.
20 Recombinant BV was generated through co-transfection using the BaculoGold
Transfection reagent (PharMingen). Recombinant BV was plaque purified and
verified through Western analysis. For protein production, SF-9 insect cells were
grown in SF-900-II medium at 2 x 10⁶/ml, and were infected at MOI of 0.5.
Purification of the His-tagged kinase used in screening was analogous to that
25 described for KDR.

Human Flt-1 Tyrosine Kinase Production and Purification

The baculoviral expression vector pVL1393 (Phar Mingen, Los Angeles,
CA) was used. A nucleotide sequence encoding poly-His₆ was placed 5' to the
30 nucleotide region encoding the entire intracellular kinase domain of human Flt-1
(amino acids 786-1338). The nucleotide sequence encoding the kinase domain was

-77-

generated through PCR using cDNA libraries isolated from HUVEC cells. The histidine residues enabled affinity purification of the protein as a manner analogous to that for KDR and ZAP70. SF-9 insect cells were infected at a 0.5 multiplicity and harvested 48 hours post infection.

5

EGFR Tyrosine Kinase Source

EGFR was purchased from Sigma (Cat # E-3641; 500 units/50 μ l) and the EGF ligand was acquired from Oncogene Research Products/Calbiochem (Cat # PF011-100).

10

Expression of ZAP70

The baculoviral expression vector used was pVL1393. (Pharmingen, Los Angeles, Ca.) The nucleotide sequence encoding amino acids M(H)6 LVPR₅S was placed 5' to the region encoding the entirety of ZAP70 (amino acids 1-619). The nucleotide sequence encoding the ZAP70 coding region was generated through PCR using cDNA libraries isolated from Jurkat immortalized T-cells. The histidine residues enabled affinity purification of the protein (vide infra). The LVPR₅S bridge constitutes a recognition sequence for proteolytic cleavage by thrombin, enabling removal of the affinity tag from the enzyme. SF-9 insect cells were infected at a multiplicity of infection of 0.5 and harvested 48 hours post infection.

20

Extraction and purification of ZAP70

SF-9 cells were lysed in a buffer consisting of 20 mM Tris, pH 8.0, 137 mM NaCl, 10% glycerol, 1% Triton X-100, 1 mM PMSF, 1 μ g/ml leupeptin, 10 μ g/ml aprotinin and 1 mM sodium orthovanadate. The soluble lysate was applied to a chelating sepharose HiTrap column (Pharmacia) equilibrated in 50 mM HEPES, pH 7.5, 0.3 M NaCl. Fusion protein was eluted with 250 mM imidazole. The enzyme was stored in buffer containing 50 mM HEPES, pH 7.5, 50 mM NaCl and 5 mM DTT.

30

-78-

Protein kinase source

Lck, Fyn, Src, Blk, Csk, and Lyn, and truncated forms thereof may be commercially obtained (e.g. from Upstate Biotechnology Inc. (Saranac Lake, N.Y) and Santa Cruz Biotechnology Inc. (Santa Cruz, Ca.)) or purified from known
5 natural or recombinant sources using conventional methods.

Enzyme Linked Immunosorbent Assay (ELISA) For PTKs

Enzyme linked immunosorbent assays (ELISA) were used to detect and measure the presence of tyrosine kinase activity. The ELISA were conducted according to known protocols which are described in, for example, Voller, *et al.*,
10 1980, "Enzyme-Linked Immunosorbent Assay," In: *Manual of Clinical Immunology*, 2d ed., edited by Rose and Friedman, pp 359-371 Am. Soc. of Microbiology, Washington, D.C.

The disclosed protocol was adapted for determining activity with respect to a specific PTK. For example, preferred protocols for conducting the ELISA
15 experiments is provided below. Adaptation of these protocols for determining a compound's activity for other members of the receptor PTK family, as well as non-receptor tyrosine kinases, are well within the abilities of those in the art. For purposes of determining inhibitor selectivity, a universal PTK substrate (e.g., random copolymer of poly(Glu₄ Tyr), 20,000-50,000 MW) was employed together
20 with ATP (typically 5 μ M) at concentrations approximately twice the apparent Km in the assay.

The following procedure was used to assay the inhibitory effect of compounds of this invention on KDR, Flt-1, Flt-4, Tie-1, Tie-2, EGFR, FGFR, PDGFR, IGF-1-R, c-Met, Lck, hck, Blk, Csk, Src, Lyn, fgr, Fyn and ZAP70 tyrosine
25 kinase activity:

Buffers and Solutions:

PGTPoly (Glu,Tyr) 4:1

Store powder at -20°C. Dissolve powder in phosphate buffered saline (PBS) for 50mg/ml solution. Store 1ml aliquots at -20°C. When making plates dilute to
30 250 μ g/ml in Gibco PBS.

Reaction Buffer: 100mM Hepes, 20mM MgCl₂, 4mM MnCl₂, 5mM DTT,

-79-

0.02%BSA, 200 μ M NaVO₄, pH 7.10

ATP: Store aliquots of 100mM at -20°C. Dilute to 20 μ M in water

Washing Buffer: PBS with 0.1% Tween 20

Antibody Diluting Buffer: 0.1% bovine serum albumin (BSA) in PBS

- 5 TMB Substrate: mix TMB substrate and Peroxide solutions 9:1 just before use or use K-Blue Substrate from Neogen

Stop Solution: 1M Phosphoric Acid

Procedure

- 10 1. Plate Preparation:

Dilute PGT stock (50mg/ml, frozen) in PBS to a 250 μ g/ml. Add 125 μ l per well of Corning modified flat bottom high affinity ELISA plates (Corning #25805-96). Add 125 μ l PBS to blank wells. Cover with sealing tape and incubate overnight 37°C.

Wash 1x with 250 μ l washing buffer and dry for about 2hrs in 37°C dry incubator.

- 15 Store coated plates in sealed bag at 4°C until used.

2. Tyrosine Kinase Reaction:

-Prepare inhibitor solutions at a 4x concentration in 20% DMSO in water.

-Prepare reaction buffer

- 20 -Prepare enzyme solution so that desired units are in 50 μ l, e.g. for KDR make to 1 ng/ μ l for a total of 50ng per well in the reactions. Store on ice.

-Make 4x ATP solution to 20 μ M from 100mM stock in water. Store on ice

-Add 50 μ l of the enzyme solution per well (typically 5-50 ng enzyme/well depending on the specific activity of the kinase)

-Add 25 μ l 4x inhibitor

- 25 -Add 25 μ l 4x ATP for inhibitor assay

-Incubate for 10 minutes at room temperature

-Stop reaction by adding 50 μ l 0.05N HCl per well

-Wash plate

**Final Concentrations for Reaction: 5 μ M ATP, 5% DMSO

- 30 3. Antibody Binding

-80-

-Dilute 1mg/ml aliquot of PY20-HRP (Pierce) antibody(a phosphotyrosine antibody)to 50ng/ml in 0.1% BSA in PBS by a 2 step dilution (100x, then 200x)

-Add 100µl Ab per well. Incubate 1 hr at room temp. Incubate 1hr at 4C.

-Wash 4x plate

5 4. Color reaction

-Prepare TMB substrate and add 100µl per well

-Monitor OD at 650nm until 0.6 is reached

-Stop with 1M Phosphoric acid. Shake on plate reader.

-Read OD immediately at 450nm

10 Optimal incubation times and enzyme reaction conditions vary slightly with enzyme preparations and are determined empirically for each lot.

For Lck, the Reaction Buffer utilized was 100 mM MOPSO, pH 6.5, 4 mM MnCl₂, 20 mM MgCl₂, 5 mM DTT, 0.2% BSA, 200 mM NaVO₄ under the analogous assay conditions.

15 Compounds of formulas 1-109 may have therapeutic utility in the treatment of diseases involving both identified, including those not mentioned herein, and as yet unidentified protein tyrosine kinases which are inhibited by compounds of formulas 1-109.

20 Cdc2 source

The human recombinant enzyme and assay buffer may be obtained commercially (New England Biolabs, Beverly, MA. USA) or purified from known natural or recombinant sources using conventional methods.

25 Cdc2 Assay

A protocol that can be used is that provided with the purchased reagents with minor modifications. In brief, the reaction is carried out in a buffer consisting of 50mM Tris pH 7.5, 100mM NaCl, 1mM EGTA, 2mM DTT, 0.01% Brij, 5% DMSO and 10mM MgCl₂ (commercial buffer) supplemented with fresh 300 µM ATP (31
30 µCi/ml) and 30 µg/ml histone type IIIss final concentrations. A reaction volume of 80µL, containing units of enzyme, is run for 20 minutes at 25 degrees C in the

-81-

presence or absence of inhibitor. The reaction is terminated by the addition of 120 μ L of 10% acetic acid. The substrate is separated from unincorporated label by spotting the mixture on phosphocellulose paper, followed by 3 washes of 5 minutes each with 75mM phosphoric acid. Counts are measured by a betacounter in the presence of liquid scintillant.

PKC kinase source

The catalytic subunit of PKC may be obtained commercially (Calbiochem).

10 PKC kinase assay

A radioactive kinase assay is employed following a published procedure (Yasuda, I., Kirshimoto, A., Tanaka, S., Tominaga, M., Sakurai, A., Nishizuka, Y. *Biochemical and Biophysical Research Communication* 3:166, 1220-1227 (1990)). Briefly, all reactions are performed in a kinase buffer consisting of 50 mM Tris-HCl pH7.5, 10mM MgCl₂, 2mM DTT, 1mM EGTA, 100 μ M ATP, 8 μ M peptide, 5% DMSO and ³³P ATP (8Ci/mM). Compound and enzyme are mixed in the reaction vessel and the reaction is initiated by addition of the ATP and substrate mixture. Following termination of the reaction by the addition of 10 μ L stop buffer (5 mM ATP in 75mM phosphoric acid), a portion of the mixture is spotted on phosphocellulose filters. The spotted samples are washed 3 times in 75 mM phosphoric acid at room temperature for 5 to 15 minutes. Incorporation of radiolabel is quantified by liquid scintillation counting.

Erk2 enzyme source

25 The recombinant murine enzyme and assay buffer may be obtained commercially (New England Biolabs, Beverly MA. USA) or purified from known natural or recombinant sources using conventional methods.

Erk2 enzyme assay

30 In brief, the reaction is carried out in a buffer consisting of 50 mM Tris pH 7.5, 1mM EGTA, 2mM DTT, 0.01% Brij, 5% DMSO and 10 mM MgCl₂

-82-

(commercial buffer) supplemented with fresh 100 μ M ATP (31 μ Ci/ml) and 30 μ M myelin basic protein under conditions recommended by the supplier. Reaction volumes and method of assaying incorporated radioactivity are as described for the PKC assay (*vide supra*).

5

In Vitro Models for T-cell Activation

Upon activation by mitogen or antigen, T-cells are induced to secrete IL-2, a growth factor that supports their subsequent proliferative phase. Therefore, one may measure either production of IL-2 from or cell proliferation of, primary T-cells or
10 appropriate T-cell lines as a surrogate for T-cell activation. Both of these assays are well described in the literature and their parameters well documented (in Current Protocols in Immunology, Vol 2, 7.10.1-7.11.2).

In brief, T-cells may be activated by co-culture with allogenic stimulator cells, a process termed the one-way mixed lymphocyte reaction. Responder and
15 stimulator peripheral blood mononuclear cells are purified by Ficoll-Hypaque gradient (Pharmacia) per directions of the manufacturer. Stimulator cells are mitotically inactivated by treatment with mitomycin C (Sigma) or gamma irradiation. Responder and stimulator cells are co-cultured at a ratio of two to one in the presence or absence of the test compound. Typically 10^5 responders are mixed
20 with 5×10^4 stimulators and plated (200 μ l volume) in a U bottom microtiter plate (Costar Scientific). The cells are cultured in RPMI 1640 supplemented with either heat inactivated fetal bovine serum (Hyclone Laboratories) or pooled human AB serum from male donors, 5×10^{-5} M 2mercaptoethanol and 0.5% DMSO. The cultures are pulsed with 0.5 μ Ci of 3 H thymidine (Amersham) one day prior to
25 harvest (typically day three). The cultures are harvested (Betaplate harvester, Wallac) and isotope uptake assessed by liquid scintillation (Betaplate, Wallac).

The same culture system may be used for assessing T-cell activation by measurement of IL-2 production. Eighteen to twenty-four hours after culture initiation, the supernatants are removed and the IL-2 concentration is measured by
30 ELISA (R and D Systems) following the directions of the manufacturer.

In-vivo Models of T-Cell Activation

The *in vivo* efficacy of compounds can be tested in animal models known to directly measure T-cell activation or for which T-cells have been proven the effectors. T-cells can be activated *in vivo* by ligation of the constant portion of the T-cell receptor with a monoclonal anti-CD3 antibody (Ab). In this model, BALB/c mice are given 10µg of anti-CD3 Ab intraperitoneally two hours prior to exsanguination. Animals to receive a test drug are pre-treated with a single dose of the compound one hour prior to anti-CD3 Ab administration. Serum levels of the proinflammatory cytokines interferon-γ (IFN- γ) and tumor necrosis factor-α(TNF-α), indicators of T-cell activation, are measured by ELISA. A similar model employs *in vivo* T-cell priming with a specific antigen such as keyhole limpet hemocyanin (KLH) followed by a secondary *in vitro* challenge of draining lymph node cells with the same antigen. As previously, measurement of cytokine production is used to assess the activation state of the cultured cells. Briefly, C57BL/6 mice are immunized subcutaneously with 100 µg KLH emulsified in complete Freund's adjuvant (CFA) on day zero. Animals are pre-treated with the compound one day prior to immunization and subsequently on days one, two and three post immunization. Draining lymph nodes are harvested on day 4 and their cells cultured at 6×10^6 per ml in tissue culture medium (RPMI 1640 supplemented with heat inactivated fetal bovine serum (Hyclone Laboratories) 5×10^{-5} M 2-mercaptoethanol and 0.5% DMSO) for both twenty-four and forty-eight hours. Culture supernatants are then assessed for the autocrine T-cell growth factor Interleukin-2 (IL-2) and/or IFN-γ levels by ELISA.

Lead compounds can also be tested in animal models of human disease. These are exemplified by experimental auto-immune encephalomyelitis (EAE) and collagen-induced arthritis (CIA). EAE models which mimic aspects of human multiple sclerosis have been described in both rats and mice (reviewed FASEB J. 5:2560-2566, 1991; murine model: Lab. Invest. 4(3):278, 1981; rodent model: J. Immunol 146(4):1163-8, 1991). Briefly, mice or rats are immunized with an emulsion of myelin basic protein (MBP), or neurogenic peptide derivatives thereof, and CFA. Acute disease can be induced with the addition of bacterial toxins

-84-

such as *bordetella pertussis*. Relapsing/remitting disease is induced by adoptive transfer of T-cells from MBP/ peptide immunized animals.

CIA may be induced in DBA/1 mice by immunization with type II collagen (J. Immunol:142(7):2237-2243). Mice will develop signs of arthritis as early as ten
5 days following antigen challenge and may be scored for as long as ninety days after immunization. In both the EAE and CIA models, a compound may be administered either prophylactically or at the time of disease onset. Efficacious drugs should reduce severity and/or incidence.

Certain compounds of this invention which inhibit one or more angiogenic
10 receptor PTK, and/or a protein kinase such as lck involved in mediating inflammatory responses can reduce the severity and incidence of arthritis in these models.

Compounds can also be tested in mouse allograft models, either skin (reviewed in Ann. Rev. Immunol., 10:333-58, 1992; Transplantation: 57(12):
15 1701-17D6, 1994) or heart (Am.J.Anat.:113:273, 1963). Briefly, full thickness skin grafts are transplanted from C57BL/6 mice to BALB/c mice. The grafts can be examined daily, beginning at day six, for evidence of rejection. In the mouse neonatal heart transplant model, neonatal hearts are ectopically transplanted from C57BL/6 mice into the ear pinnae of adult CBA/J mice. Hearts start to beat four to
20 seven days post transplantation and rejection may be assessed visually using a dissecting microscope to look for cessation of beating.

Cellular Receptor PTK Assays

The following cellular assay was used to determine the level of activity and
25 effect of the different compounds of the present invention on KDR/VEGFR2. Similar receptor PTK assays employing a specific ligand stimulus can be designed along the same lines for other tyrosine kinases using techniques well known in the art.

30 VEGF-Induced KDR Phosphorylation in Human Umbilical Vein Endothelial Cells (HUVEC) as Measured by Western Blots:

-85-

1. HUVEC cells (from pooled donors) can be purchased from Clonetics (San Diego, CA) and cultured according to the manufacturer directions. Only early passages (3-8) are used for this assay. Cells are cultured in 100 mm dishes (Falcon for tissue culture; Becton Dickinson; Plymouth, England) using complete EBM media (Clonetics).

2. For evaluating a compound's inhibitory activity, cells are trypsinized and seeded at $0.5-1.0 \times 10^5$ cells/well in each well of 6-well cluster plates (Costar; Cambridge, MA).

3. 3-4 days after seeding, plates are typically 90-100% confluent.
10 Medium is removed from all the wells, cells are rinsed with 5-10ml of PBS and incubated 18-24h with 5ml of EBM base media with no supplements added (i.e., serum starvation).

4. Serial dilutions of inhibitors are added in 1ml of EBM media (25 μ M, 5 μ M, or 1 μ M final concentration to cells and incubated for one hour at 37 C.
15 Human recombinant VEGF₁₆₅ (R & D Systems) is then added to all the wells in 2 ml of EBM medium at a final concentration of 50ng/ml and incubated at 37 C for 10 minutes. Control cells untreated or treated with VEGF only are used to assess background phosphorylation and phosphorylation induction by VEGF.

All wells are then rinsed with 5-10ml of cold PBS containing 1mM Sodium
20 Orthovanadate (Sigma) and cells are lysed and scraped in 200 μ l of RIPA buffer (50mM Tris-HCl) pH7, 150mM NaCl, 1% NP-40, 0.25% sodium deoxycholate, 1mM EDTA) containing protease inhibitors (PMSF 1mM, aprotinin 1 μ g/ml, pepstatin 1 μ g/ml, leupeptin 1 μ g/ml, Na vanadate 1mM, Na fluoride 1mM) and 1 μ g/ml of Dnase (all chemicals from Sigma Chemical Company, St Louis, MO).
25 The lysate is spun at 14,000 rpm for 30min, to eliminate nuclei.

Equal amounts of proteins are then precipitated by addition of cold (-20 C) Ethanol (2 volumes) for a minimum of 1 hour or a maximum of overnight. Pellets are reconstituted in Laemli sample buffer containing 5% β -mercaptoethanol (BioRad; Hercules, CA) and boiled for 5min. The proteins are resolved by polyacrylamide gel
30 electrophoresis (6%, 1.5mm Novex, San Deigo, CA) and transferred onto a nitrocellulose membrane using the Novex system. After blocking with bovine serum

-86-

albumin (3%), the proteins are probed overnight with anti-KDR polyclonal antibody (C20, Santa Cruz Biotechnology; Santa Cruz, CA) or with anti-phosphotyrosine monoclonal antibody (4G10, Upstate Biotechnology, Lake Placid, NY) at 4 C. After washing and incubating for 1 hour with HRP-conjugated F(ab)₂ of goat anti-rabbit or
5 goat-anti-mouse IgG the bands are visualized using the emission chemiluminescence (ECL) system (Amersham Life Sciences, Arlington Height, IL).

In vivo Uterine Edema Model

This assay measures the capacity of compounds to inhibit the acute increase
10 in uterine weight in mice which occurs in the first few hours following estrogen stimulation. This early onset of uterine weight increase is known to be due to edema caused by increased permeability of uterine vasculature. Cullinan-Bove and Koss (*Endocrinology* (1993), 133:829-837) demonstrated a close temporal relationship of estrogen-stimulated uterine edema with increased expression of VEGF mRNA in the
15 uterus. These results have been confirmed by the use of neutralizing monoclonal antibody to VEGF which significantly reduced the acute increase in uterine weight following estrogen stimulation (WO 97/42187). Hence, this system can serve as a model for *in vivo* inhibition of VEGF signalling and the associated hyperpermeability and edema.

20 **Materials:** All hormones can be purchased from Sigma (St. Louis, MO) or Cal Biochem (La Jolla, CA) as lyophilized powders and prepared according to supplier instructions.

Vehicle components (DMSO, Cremaphor EL) can be purchased from Sigma (St. Louis, MO).

25 **Mice** (Balb/c, 8-12 weeks old) can be purchased from Taconic (Germantown, NY) and housed in a pathogen-free animal facility in accordance with institutional Animal Care and Use Committee Guidelines.

Method:

Day 1: Balb/c mice are given an intraperitoneal (i.p.) injection of 12.5
30 units of pregnant mare's serum gonadotropin (PMSG).

Day 3: Mice receive 15 units of human chorionic gonadotropin

-87-

(hCG) i.p.

Day 4: Mice are randomized and divided into groups of 5-10. Test compounds are administered by i.p., i.v. or p.o. routes depending on solubility and vehicle at doses ranging from 1-100 mg/kg. Vehicle control group receive vehicle only and two groups are left untreated.

Thirty minutes later, experimental, vehicle and 1 of the untreated groups are given an i.p. injection of 17 -estradiol (500 µg/kg). After 2-3 hours, the animals are sacrificed by CO₂ inhalation. Following a midline incision, each uterus was isolated and removed by cutting just below the cervix and at the junctions of the uterus and oviducts. Fat and connective tissue were removed with care not to disturb the integrity of the uterus prior to weighing (wet weight). Uteri are blotted to remove fluid by pressing between two sheets of filter paper with a one liter glass bottle filled with water. Uteri are weighed following blotting (blotted weight). The difference between wet and blotted weights is taken as the fluid content of the uterus. Mean fluid content of treated groups is compared to untreated or vehicle treated groups. Significance is determined by Student's test. Non-stimulated control group is used to monitor estradiol response.

Certain compounds of this invention which are inhibitors of angiogenic receptor tyrosine kinases can also be shown active in a Matrigel implant model of neovascularization. The Matrigel neovascularization model involves the formation of new blood vessels within a clear marble of extracellular matrix implanted subcutaneously which is induced by the presence of proangiogenic factor producing tumor cells (for examples see: Passaniti, A., *et al*, Lab. Investig. (1992), 67(4), 519-528; Anat. Rec. (1997), 249(1), 63-73; Int. J. Cancer (1995), 63(5), 694-701; Vasc. Biol. (1995), 15(11), 1857-6). The model preferably runs over 3-4days and endpoints include macroscopic visual/image scoring of neovascularization, microscopic microvessel density determinations, and hemoglobin quantitation (Drabkin method) following removal of the implant versus controls from animals untreated with inhibitors. The model may alternatively employ bFGF or HGF as the stimulus.

Example 1:

7-Cyclopentyl-5-(4-phenoxyphenyl)-5*H*-pyrrolo[3,2-*d*]pyrimidin-4-amine

5 a) 2-cyclopentylacetonitrile

A mixture of sodium hydride (2.17 g, 60% in oil, 54.2 mmol) in diethyl ether (100 mL) was cooled to 0°C then treated with diethyl(cyanomethyl) phosphonate (9.6 g, 54.2 mmol) while maintaining the temperature of the mixture at less than 0°C.

Cyclopentanone (4.13 g, 49.3 mmol) in diethyl ether (25 mL) was added to the
10 mixture at less than 5°C then the reaction was warmed to ambient temperature and stirred for an additional 16 hours. Water (240 mL) was added to the mixture and the layers were then separated. The aqueous layer was extracted with diethyl ether (50 mL). The combined organic solutions were extracted with water (50 mL) then brine (50 mL) and finally dried over magnesium sulfate, filtered and the filtrate
15 concentrated under reduced pressure. The resulting residue was dissolved in ethanol (40 mL) then 10% palladium on carbon (250 mg) was added and the mixture hydrogenated at atmospheric pressure and ambient temperature for 16 hours. The catalyst was removed by filtration through a pad of celite and the filtrate concentrated to an oil under reduced pressure. The title compound was purified by
20 fractional distillation to give 4.08 g (75.6%) as a light yellow oil (boiling point 63°C at 20 torr). : ¹H NMR (Chloroform-*d*, 400 MHz) δ 2.35 (d, 2H), 2.18 (m, 1H), 1.87 (m, 2H), 1.59-1.69 (m, 4H), 1.29 (m, 2H).

b) 1-cyclopentyl-2-oxoethylcyanide

25 A mixture of 2-cyclopentylacetonitrile (0.50 g, 4.59 mmol) in tetrahydrofuran (10 mL) was cooled to -60°C then treated with 1.7 M *tert*-butyllithium in pentane (3.25 mL, 5.50 mmol) while maintaining the reaction temperature at less than -55°C. The solution was stirred for 10 minutes then ethyl formate (0.41 g, 5.50 mmol) was added dropwise. The mixture was warmed to ambient temperature and stirred an
30 additional 16 hours. The mixture was concentrated under reduced pressure and the resulting residue applied to a silica gel column and eluted with dichloromethane/ethyl acetate (95:5). The fractions containing material with an R_f

of 0.1-0.3 [TLC, dichloromethane/ethyl acetate (95:5), potassium permanganate stain) were combined and concentrated to give an oil which was used without further purification: ^1H NMR (Chloroform- d , 400 MHz) δ 9.57 (s, 1H), 3.54 (d, 1H), 2.45 (m, 1H), 1.4-1.9 (m, 8H).

5

c) (4-phenoxyanilino)methyl cyanide

A mixture of 4-phenoxyaniline (7.0 g, 37.8 mmol), bromoacetonitrile (4.5 g, 37.8 mmol) and triethylamine (4.2 g, 41.6 mmol) in tetrahydrofuran (50 mL) was heated at 85°C for 5.25 hours then cooled and another portion of bromoacetonitrile (6.5 g, 5.46 mmol) was added. The mixture was heated at 85°C for 18 hours then cooled and concentrated under reduced pressure. The residue was partitioned between dichloromethane (50 mL) and water (50 mL). The aqueous layer was extracted with dichloromethane (30 mL) then the combined organic solutions were extracted with 5 N aqueous sodium hydroxide (30 mL), dried over magnesium sulfate, filtered and the filtrate concentrated under reduced pressure. The residue was then purified by flash chromatography on silica gel using dichloromethane/ethyl acetate (98:2) as an eluent to provide 3.8 g (45%) of the title compound as a dark brown solid: ^1H NMR (DMSO- d_6 , 400 MHz) δ 7.32 (t, 2H), 7.03 (t, 1H), 6.87-6.94 (m, 4H), 6.76 (d, 2H), 6.26 (bs, 2H), 4.25 (s, 2H); RP-HPLC (Hypersil HS-C18, 5 μm , 100A, 4.6 x 250 mm; 25%-100% acetonitrile-0.05 M ammonium acetate over 25 min, 1 mL/min) t_r 18.4 min.; MS: MH^+ 225.1.

d) 3-amino-4-cyclopentyl-1-(4-phenoxyphenyl)-1*H*-2-pyrrolicarbonitrile

A mixture of (4-phenoxyanilino)methyl cyanide (0.68 g, 3.30 mmol) and 1-cyclopentyl-2-oxoethylcyanide (0.54 g, 3.94 mmol) in 1,2-dimethoxyethane (10 mL) was treated with 2 drops of acetic acid then heated at 85°C for 45 minutes. The mixture was cooled to ambient temperature then 1,5-diazabicyclo[4.3.0]non-5-ene (DBN) (1.13 g, 9.09 mmol) was added. The mixture was then heated at 65°C for 16 hours and 85°C for 6 hours. Fresh DBN (0.25 mL) was added and the mixture was heated at 85°C for an additional 18 hours. The solvent was removed under reduced pressure then the residue was applied to a silica gel column and eluted with

-90-

heptane/ethyl acetate (7:3) to provide 185 mg (17.8%) of the title compound as a glass: ¹H NMR (DMSO-*d*₆, 400 MHz) δ 7.42 (m, 4H), 7.17 (t, 1H), 7.04-7.11 (m, 4H), 6.99 (s, 1H), 5.10 (bs, 2H), 2.82 (m, 1H), 1.97 (m, 2H), 1.69 (m, 2H), 1.58 (m, 2H); RP-HPLC (Hypersil HS-C18, 5μm, 100A, 4.6 x 250 mm; 25%-100% acetonitrile-0.05 M ammonium acetate over 25 min, 1 mL/min) *t*_r 26.2 min.; MS: MH⁺ 343.9.

e) 7-cyclopentyl-5-(4-phenoxyphenyl)-5*H*-pyrrolo[3,2-*d*]pyrimidin-4-amine
A mixture of the 3-amino-4-cyclopentyl-1-(4-phenoxyphenyl)-1*H*-2-pyrrolocarbonitrile (185 mg, 0.539 mmol) in absolute ethanol (10 mL) was treated with formamidine acetate (450 mg, 4.33 mmol) then heated at 85°C for 2 hours. The solvent was evaporated under reduced pressure then the residue was purified by preparative reverse phase HPLC to provide 145 mg (73%) of the title compound as a white solid after lyophilization: ¹H NMR (DMSO-*d*₆, 400 MHz) δ 8.19 (s, 1H), 7.44 (m, 5H), 7.19 (t, 1H), 7.13 (m, 4H), 5.79 (bs, 2H), 3.23 (m, 1H), 2.05 (m, 2H), 1.77 (m, 4H), 1.64 (m, 2H); RP-HPLC (Hypersil HS-C18, 5μm, 100A, 4.6 x 250 mm; 5%-100% acetonitrile-0.05 M ammonium acetate over 25 min, 1 mL/min) *t*_r 23.3min.; MS: MH⁺ 371.5.

20 Example 2:

1-cyclopentyl-3-(4-phenoxyphenyl)-1*H*-pyrrolo-[2,3-*d*] pyridazin-4-amine

a) Ethyl 3-cyano-1-cyclopentyl-1*H*-2-pyrrolocarboxylate

The title compound was prepared from cyclopentyl amine and diethyl (2*E*, 4*E*, 6*E*)-3,6-dicyano-2,7-dihydroxy-2,4,6-octatriendioate in a 13% yield by the method described by Huisgen¹: ¹H NMR (DMSO-*d*₆, 400 MHz) δ 7.47 (d, 1H), 6.72 (d, 1H), 5.37 (m, 1H), 4.31 (q, 2H), 2.11 (m, 2H), 1.77 (m, 4H), 1.66 (m, 2H), 1.32 (t, 3H); RP-HPLC (Hypersil HS-C18, 5μm, 100A, 4.6 x 250 mm; 5%-100% acetonitrile-0.05 M ammonium acetate over 25 min, 1 mL/min) *t*_r 22.2 min.

30

b) 3-cyano-1-cyclopentyl-1*H*-2-pyrrolocarboxylic acid

-91-

A mixture of ethyl 3-cyano-1-cyclopentyl-1*H*-2-pyrrolocarboxylate (1.15 g, 5.16 mmol) in ethanol (25 mL) and water (5 mL) was treated with potassium hydroxide (0.58 g, 10.32 mmol). The mixture was heated at 75°C for 30 minutes then cooled and the solvents evaporated under reduced pressure. Water (20 mL) was added and the solution was cooled to 0°C then acidified with concentrated hydrochloric acid (1.3 g, 36% by weight, 11.35 mmol). The slurry that formed was stirred for 10 minutes then the solid was collected by filtration to give the title compound (0.75 g, 75%) as a light orange solid. ¹H NMR (DMSO-*d*₆, 400 MHz) δ 13.49 (bs, 1H), 7.41 (d, 1H), 6.67 (d, 1H), 5.47 (m, 1H), 2.20 (m, 2H), 1.77 (m, 4H), 1.64 (m, 2H); RP-HPLC (Hypersil HS-C18, 5μm, 100A, 4.6 x 250 mm; 5%-100% acetonitrile-0.05 M ammonium acetate over 25 min, 1 mL/min) *t*_r 12.52 min.

c) 1-cyclopentyl-2-formyl-1*H*-3-pyrrolocarbonitrile

A mixture of 3-cyano-1-cyclopentyl-1*H*-2-pyrrolocarboxylic acid (0.75 g, 3.68 mmol) in dichloromethane (5 mL) was cooled to 0°C then treated with oxalyl chloride (0.52 g, 4.04 mmol). *N,N*-Dimethylformamide (1 drop) was added then the mixture was warmed to ambient temperature and stirred for 1.5 hours. The solvents were removed under reduced pressure then the residue was dissolved in diglyme (10 mL). The solution which resulted was cooled to -60°C then lithium tri-*tert*-butoxyaluminumhydride (8 mL, 0.5 M solution in diglyme, 4.0 mmol) was added dropwise over the course of approximately 1.5 hours while maintaining the temperature of the solution below -60°C. The mixture was allowed to warm to -10°C then it was cooled to -60°C and an additional portion of 0.5 M lithium tri-*tert*-butoxyaluminumhydride in diglyme was added (1.5 mL, 0.75 mmol). The mixture was warmed to ambient temperature then treated with concentrated hydrochloric acid (1 mL) and purified by preparative reverse phase HPLC to provide the title compound (200 mg, 30%): ¹H NMR (DMSO-*d*₆, 400 MHz) δ 9.82 (s, 1H), 7.60 (d, 1H), 6.84 (d, 1H), 5.32 (m, 2H), 2.10 (m, 2H), 1.81 (m, 4H), 1.66 (m, 2H); 5%-100% acetonitrile-0.05 M ammonium acetate over 25 min, 1 mL/min) *t*_r 19.46 min. *m/z* GC/MS: MH⁺ 189.2.

-92-

d) 1-cyclopentyl-1*H*-pyrrolo[2,3-*d*]pyridazin-4-amine

A mixture of 1-cyclopentyl-2-formyl-1*H*-3-pyrrolicarbonitrile (0.525 g, 2.79 mmol) and hydrazine dihydrochloride (0.35 g, 3.35 mmol) in ethanol (30 mL) was heated at reflux for 2.5 hours then cooled to ambient temperature and purified by preparative reverse phase HPLC to give the title compound as a hygroscopic glass contaminated with ammonium acetate (594 mg) (60% by weight as determined by ¹H NMR, yield= 337 mg; ¹H NMR (DMSO-*d*₆, 400 MHz) δ 8.83 (s, 1H), 7.55 (d, 1H), 6.71 (d, 1H), 6.10 (bs, 2H), 4.94 (m, 1H), 2.15 (m, 2H), 1.89 (m, 4H), 1.70 (m, 2H); 5%-100% acetonitrile-0.05 M ammonium acetate over 25 min, 1 mL/min) *t*_r 19.46 min. .; LC/MS: MH⁺ 203.0.

e) 3-bromo-1-cyclopentyl-1*H*-pyrrolo[2,3-*d*]pyridazin-4-amine

A mixture of 1-cyclopentyl-1*H*-pyrrolo[2,3-*d*]pyridazin-4-amine (0.595 mg, approx. 60% pure, 1.76 mmol) in dichloromethane (100 mL) was treated with a solution of dichloromethane (5 mL) containing bromine (0.5 g, 2.95 mmol) over the course of 1.25 hours. The mixture was stirred an additional one hour then another portion of dichloromethane (3 mL) containing bromine (0.3 g) was added. The mixture was stirred for 2.5 hours then treated with 5 mL 5N aqueous sodium hydroxide and 25 mL water. The layers were separated and the organic layer was concentrated under reduced pressure to give a residue which was purified by reverse phase preparative HPLC to give 3-bromo-1-cyclopentyl-1*H*-pyrrolo[2,3-*d*]pyridazin-4-amine (168 mg, 35%; ¹H NMR (DMSO-*d*₆, 400 MHz) δ 8.94 (s, 1H), 7.79 (s, 1H), 6.11 (bs, 2H), 4.94 (m, 1H), 2.12 (m, 2H), 1.83 (m, 4H), 1.67 (m, 2H); 5%-100% acetonitrile-0.05 M ammonium acetate over 25 min, 1 mL/min) *t*_r 11.25 min. .; LC/MS: MH⁺ 282.8.

f) 1-cyclopentyl-3-(4-phenoxyphenyl)-1*H*-pyrrolo-[2,3-*d*] pyridazin-4-amine


A mixture of 3-bromo-1-cyclopentyl-1*H*-pyrrolo[2,3-*d*]pyridazin-4-amine (0.057 g, 0.178 mmol), 4-phenoxyphenyl boronic acid (0.057g, 0.266 mmol), sodium carbonate (0.062 g, 0.588 mmol) and tetrakis(triphenylphosphine)palladium(0) (12 mg, 0.011 mmol) in ethylene glycol

-93-

dimethyl ether (3 mL) and water (1.5 mL) was heated at 85°C under an atmosphere of nitrogen for 2.5 hours. The mixture was cooled to ambient temperature and the solvent evaporated under reduced pressure. The residue was purified by preparative reverse phase HPLC to give the title compound contaminated with 4-phenoxyphenyl boronic acid. The residue was partitioned between dichloromethane and 5 N aqueous sodium hydroxide. The organic layer was dried over magnesium sulfate, filtered and the filtrate evaporated to give the title compound (14 mg, 21%); ¹H NMR (DMSO-*d*₆, 400 MHz) δ 8.98 (s, 1H), 7.68 (s, 1H), 7.50 (d, 2H), 7.40 (t, 2H), 7.17 (t, 1H), 7.09 (m, 4H), 5.76 (bs, 2H), 5.01 (m, 1H), 2.19 (m, 2H), 1.91 (m, 2H), 1.71 (m, 2H); 5%-100% acetonitrile-0.05 M ammonium acetate over 25 min, 1 mL/min) t_r 20.60 min. ; LC/MS: MH⁺ 371.2.

(1) Huisgen, R.; Laschtuvka, E. Eine Neue Synthesen von Derivaten des Pyrroles. Chem. Ber. 1960, 93, 65.

Other preferred compounds of the instant invention are those compounds where the compound of Formula (I) is



R₁ = trans-2-phenyl-cyclopropanecarboxamide
 phenyl-4-(trifluoromethyl)benzamide
 phenyl-1-methyl-2-indolecarboxamide
 phenyl-2,2-dimethyl-3-phenylpropanamide
 phenoxyphenyl

R₂ = cyclopentyl
 cis-cyclohexyl piperazine
 trans-cyclohexyl piperazine
 piperaziny-piperiziny

More specifically, the compounds are:

*N*1-[4-(7-amino-3-cyclopentyl-1*H*-pyrazolo[3,4-*c*]pyridin-1-yl)phenyl]-*trans*-2-phenylcyclopropane-1-carboxamide;

- Cis-N1*-(4-{7-amino-3-[4-(4-methylpiperazino)cyclohexyl]-1*H*-pyrazolo[3,4-*c*]pyridin-1-yl}phenyl)-*trans*-2-phenylcyclopropane-1-carboxamide;
- Trans-N1*-(4-{7-amino-3-[4-(4-methylpiperazino)cyclohexyl]-1*H*-pyrazolo[3,4-*c*]pyridin-1-yl}phenyl)-*trans*-2-phenylcyclopropane-1-carboxamide;
- 5 *N1*-(4-{7-amino-3-(1-methylpiperidin-4-yl)piperidin-4-yl]-1*H*-pyrazolo[3,4-*c*]pyridin-1-yl}phenyl)-2-phenyl-1-cyclopropanecarboxamide;
- N1*-[4-(7-amino-3-cyclopentyl-1*H*-pyrazolo[3,4-*c*]pyridin-1-yl)phenyl]-4-(trifluoromethyl)benzamide;
- Cis-N1*-(4-{7-amino-3-[4-(4-methylpiperazino)cyclohexyl]-1*H*-pyrazolo[3,4-*c*]pyridin-1-yl}phenyl)-4-(trifluoromethyl)benzamide;
- 10 *Trans-N1*-(4-{7-amino-3-[4-(4-methylpiperazino)cyclohexyl]-1*H*-pyrazolo[3,4-*c*]pyridin-1-yl}phenyl)-4-(trifluoromethyl)benzamide;
- N1*-(4-{7-amino-3-(1-methylpiperidin-4-yl)piperidin-4-yl]-1*H*-pyrazolo[3,4-*c*]pyridin-1-yl}phenyl)-4-(trifluoromethyl)benzamide;
- 15 *N2*-[4-(7-amino-3-cyclopentyl-1*H*-pyrazolo[3,4-*c*]pyridin-1-yl)phenyl]-1-methyl-1*H*-2-indolecarboxamide;
- Cis-N2*-(4-{7-amino-3-[4-(4-methylpiperazino)cyclohexyl]-1*H*-pyrazolo[3,4-*c*]pyridin-1-yl}phenyl)-1-methyl-1*H*-2-indolecarboxamide;
- Trans-N2*-(4-{7-amino-3-[4-(4-methylpiperazino)cyclohexyl]-1*H*-pyrazolo[3,4-*c*]pyridin-1-yl}phenyl)-1-methyl-1*H*-2-indolecarboxamide;
- 20 *N2*-(4-{7-amino-3-[(1-methylpiperidin-4-yl)piperidin-4-yl]-1*H*-pyrazolo[3,4-*c*]pyridin-1-yl}phenyl)-1-methyl-1*H*-2-indolecarboxamide;
- N1*-[4-(7-amino-3-cyclopentyl-1*H*-pyrazolo[3,4-*c*]pyridin-1-yl)phenyl]-2,2-dimethyl-3-phenylpropanamide;
- 25 *Cis-N1*-(4-{7-amino-3-[4-(4-methylpiperazino)cyclohexyl]-1*H*-pyrazolo[3,4-*c*]pyridin-1-yl}phenyl)-2,2-dimethyl-3-phenylpropanamide;
- Trans-N1*-(4-{7-amino-3-[4-(4-methylpiperazino)cyclohexyl]-1*H*-pyrazolo[3,4-*c*]pyridin-1-yl}phenyl)-2,2-dimethyl-3-phenylpropanamide;
- N1*-(4-{7-amino-3-[(1-methylpiperidin-4-yl)piperidin-4-yl]-1*H*-pyrazolo[3,4-*c*]pyridin-1-yl}phenyl)-2,2-dimethyl-3-phenylpropanamide;
- 30 3-cyclopentyl-1-(4-phenoxyphenyl)-1*H*-pyrazolo[3,4-*c*]pyridin-7-amine;
- Cis*-3-[4-(4-methylpiperazino)cyclohexyl]-1-(4-phenoxyphenyl)-1*H*-pyrazolo[3,4-

-95-

c]pyridin-7-amine;

Trans-3-[4-(4-methylpiperazino)cyclohexyl]-1-(4-phenoxyphenyl)-1*H*-pyrazolo[3,4-
c]pyridin-7-amine; and

3-[(1-methylpiperidin-4-yl)piperidin-4-yl]-1-(4-phenoxyphenyl)-1*H*-pyrazolo[3,4-
5 c]pyridin-7-amine.

Other preferred compounds include:

7-cyclopentyl-5-(4-phenoxyphenyl)-5*H*-pyrrolo[3,2-*d*]pyrimidin-4-amine;

1-cyclopentyl-3-(4-phenoxyphenyl)-1*H*-pyrrolo[2,3-*d*]pyridazin-4-amine;

4-amino-9-cyclopentyl-6-(4-phenoxyphenyl)-6,7,8,9-tetrahydro-5*H*-pyrimido[4,5-

10 b][1,4]diazepin-8-one; and

4-amino-9-cyclopentyl-6-(4-phenoxyphenyl)-8,9-dihydro-5*H*-pyrimido[4,5-

b][1,4]diazepin-8-one.


The foregoing compounds can be synthesized substantially according to
Examples 1 or 2 using the appropriate starting materials.

15


CLAIMS



We claim:

1. A compound of Formula (I), the racemic-diastereomeric mixtures, optical isomers, pharmaceutically-acceptable salts, prodrugs or biologically active metabolites thereof, selected from the group consisting of





-97-










5



-99-




5




10

-100-




-101-




85




86




87




88




89




90




91




92




93




94




95




96




97



98




99




100




101



102




103




104


-102-




105




106




107




108




109




110




111




112




113



114




115




116

and




117


wherein:

R₁ is

-103-



where Z^{100} is $M_2=L_2$ or a group optionally substituted with R_b ,
 selected from the group consisting of cycloalkyl, naphthyl,
 tetrahydronaphthyl, benzothienyl, furanyl, thienyl, benzoxazolyl,



benzothiazolyl, , thiazolyl, benzofuranyl,
 2,3-dihydrobenzofuranyl, indolyl, isoxazolyl, tetrahydropyranyl,
 tetrahydrofuranyl, piperidinyl, pyrazolyl, pyrrolyl, oxazolyl, isothiazolyl,
 oxadiazolyl, thiadiazolyl, indolyl, indazolyl, benzoisothiazolyl, pyrido-
 oxazolyl, pyrido-thiazolyl, pyrimido-oxazolyl, pyrimido-thiazolyl and
 benzimidazolyl;

Z^{110} is a covalent bond, or an optionally substituted (C_1-C_6) which is
 optionally substituted with one or more substituents selected from the group
 consisting of alkyl, CN, OH, halogen, NO_2 , COOH, substituted or
 unsubstituted amino and substituted or unsubstituted phenyl;

Z^{111} is a covalent bond, an optionally substituted (C_1-C_6) or an optionally
 substituted

$-(CH_2)_n$ -cycloalkyl- $(CH_2)_n$ -; where the optionally substituted groups are
 optionally substituted with one or more substituents selected from the group
 consisting of alkyl, CN, OH, halogen, NO_2 , COOH, substituted or
 unsubstituted amino and substituted or unsubstituted phenyl;

R_a and R_i each represent one or more substituents for each occurrence
 independently selected from the group consisting of hydrogen, halogen, -CN,
 - NO_2 , -C(O)OH, -C(O)H, -OH, -C(O)O-alkyl, substituted or unsubstituted
 carboxamido, tetrazolyl, trifluoromethylcarbonylamino,
 trifluoromethylsulfonamido, substituted or unsubstituted alkyl, substituted or
 unsubstituted alkoxy, substituted or unsubstituted aryl, substituted or
 unsubstituted alkenyl, substituted or unsubstituted aryloxy, substituted or

-104-

- unsubstituted heteroaryloxy, substituted or unsubstituted arylalkyl, substituted or unsubstituted alkynyl, substituted or unsubstituted amino, substituted or unsubstituted aminoalkyl, substituted or unsubstituted amido groups, substituted or unsubstituted heteroarylthio, substituted or unsubstituted arylthio, $-Z^{105}-C(O)N(R)_2$, $-Z^{105}-N(R)-C(O)-Z^{200}$, $-Z^{105}-N(R)-S(O)_2-Z^{200}$, $-Z^{105}-N(R)-C(O)-N(R)-Z^{200}$, R_e and CH_2OR_e ;
- where R_e for each occurrence is independently hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted aryl, $-CH_2-NR_dR_e$, $-W-(CH_2)_t-NR_dR_e$, $-W-(CH_2)_t-Oalkyl$, $-W-(CH_2)_t-S-alkyl$, or $-W-(CH_2)_t-OH$;
- Z^{105} for each occurrence is independently a covalent bond or (C_1-C_6) ;
- Z^{200} for each occurrence is independently a substituted or unsubstituted (C_1-C_6) , substituted or unsubstituted phenyl or substituted or unsubstituted $-(C_1-C_6)$ -phenyl;
- R_d and R_e for each occurrence are independently H, alkyl, alkanoyl or SO_2 -alkyl; or R_d , R_e and the nitrogen atom to which they are attached together form a five- or six-membered heterocyclic ring; t for each occurrence is independently an integer from 2 to 6; W for each occurrence is independently a direct bond or O, S, $S(O)$, $S(O)_2$, or NR_f , wherein R_f for each occurrence is independently H or alkyl;
- or R_i is a substituted or unsubstituted carbocyclic or heterocyclic ring fused with ring 2;
- R_3 is hydrogen, hydroxy, substituted or unsubstituted alkyl or substituted or unsubstituted alkoxy;
- A is $-O-$; $-S-$; $-S(O)_p-$; $-N(R)-$; $-N(C(O)OR)-$; $-N(C(O)R)-$; $-N(SO_2R)-$; $-CH_2O-$; $-CH_2S-$; $-CH_2N(R)-$; $-CH(NR)-$; $-CH_2N(C(O)R)-$; $-CH_2N(C(O)OR)-$; $-CH_2N(SO_2R)-$; $-CH(NHR)-$; $-CH(NHC(O)R)-$; $-CH(NHSO_2R)-$; $-CH(NHC(O)OR)-$; $-CH(OC(O)R)-$; $-CH(OC(O)NHR)-$; $-CH=CH-$; $-C(=NOR)-$; $-C(O)-$; $-CH(OR)-$; $-C(O)N(R)-$; $-N(R)C(O)-$; $-N(R)S(O)_p-$; $-OC(O)N(R)-$; $-N(R)-C(O)-(CH_2)_n-N(R)-$; $-N(R)C(O)O-$; $-N(R)-(CH_2)_{n+1}-C(O)-$; $-S(O)_pN(R)-$; $-O-(CR_2)_{n+1}-C(O)-$;

-105-

$-O-(CR_2)_{n+1}-O-$,
 $-N(C(O)R)S(O)_p-$; $-N(R)S(O)_pN(R)-$; $-N(R)-C(O)-(CH_2)_n-O-$, $-$
 $C(O)N(R)C(O)-$; $-S(O)_pN(R)C(O)-$; $-OS(O)_pN(R)-$; $-N(R)S(O)_pO-$; $-$
 $N(R)S(O)_pC(O)-$; $-SO_pN(C(O)R)-$; $-N(R)SO_pN(R)-$; $-C(O)O-$; $-$
 $N(R)P(OR_g)O-$; $-N(R)P(OR_g)-$; $-N(R)P(O)(OR_g)O-$; $-N(R)P(O)(OR_g)-$
 $;$
 $-N(C(O)R)P(OR_g)O-$; $-N(C(O)R)P(OR_g)-$; $-N(C(O)R)P(O)(OR_g)O-$,
 or
 $-N(C(O)R)P(OR_g)-$;

10 where R for each occurrence is independently H, substituted
 or unsubstituted alkyl, substituted or unsubstituted arylalkyl
 or substituted or unsubstituted aryl;

15 R_g for each occurrence is independently H, substituted or
 unsubstituted alkyl, substituted or unsubstituted arylalkyl,
 substituted or unsubstituted cycloalkyl or substituted or
 unsubstituted aryl;

p is 1 or 2;

20 or in a phosphorus containing group, the nitrogen atom, the
 phosphorus atom, R and R_g together form a five- or six-
 membered heterocyclic ring; or

A is $NRSO_2$ and R, R_a and the nitrogen atom together form a substituted or
 unsubstituted five or-six-membered heterocyclic ring fused to ring 1;

R_2 is $-Z^{101}-Z^{102}$;

25 Z^{101} is a covalent bond, $-(C_1-C_6)-$, $-(C_1-C_6)-O-$, $-(C_1-C_6)-C(O)-$, $-(C_1-C_6)-$
 $C(O)O-$, $-(C_1-C_6)-C(O)-NH-$, $-(C_1-C_6)-C(O)-N((C_1-C_6))-$ or a
 substituted or unsubstituted phenyl group;

30 Z^{102} is hydrogen, a substituted or unsubstituted alkyl group, a substituted or
 unsubstituted cycloalkyl group, a substituted or unsubstituted, saturated or
 unsaturated heterocyclic group, or a substituted or unsubstituted, saturated or
 unsaturated heterobicyclic group;

said substituted heterocyclic or substituted heterobicyclic group having one
 or more substituents each independently selected from the group consisting

of hydroxyl, cyano, substituted or unsubstituted alkoxy, substituted or unsubstituted sulfonamido, substituted or unsubstituted ureido, substituted or unsubstituted carboxamido; substituted or unsubstituted amino, oxo, a saturated, unsaturated or aromatic, substituted or unsubstituted heterocyclic group comprising one or more nitrogen atoms, one or more oxygen atoms or a combination thereof;

wherein said nitrogen atoms are independently optionally substituted by a substituted or unsubstituted alkyl, substituted or unsubstituted aryl or substituted or unsubstituted arylalkyl group; or

R_2 is of the formula B-E, wherein B is a substituted or unsubstituted cycloalkyl, substituted or unsubstituted azacycloalkyl, substituted or unsubstituted amino, substituted or unsubstituted aminoalkylsulfonyl, substituted or unsubstituted alkoxyalkyl, substituted or unsubstituted alkoxy, substituted or unsubstituted aminoalkylcarbonyl, hydroxy, substituted or unsubstituted alkylene, substituted or unsubstituted aminoalkyl, substituted or unsubstituted alkylencarbonyl or substituted or unsubstituted aminoalkylcarbonyl group; and E is substituted or unsubstituted azacycloalkyl, substituted or unsubstituted azacycloalkylcarbonyl, substituted or unsubstituted azacycloalkylsulfonyl, substituted or unsubstituted azacycloalkylalkyl, substituted or unsubstituted heteroaryl, substituted or unsubstituted heteroarylcarbonyl, substituted or unsubstituted heteroarylsulfonyl, substituted or unsubstituted heteroarylalkyl, substituted or unsubstituted azacycloalkylcarbonylamino, substituted or unsubstituted heteroarylcarbonylamino or substituted or unsubstituted aryl;

a is 1 and D_1 , G_1 , J_1 , L_1 and M_1 are each independently selected from the group

consisting of CR_a and N, provided that at least two of D_1 , G_1 , J_1 , L_1 and M_1 are CR_a ; or

a is 0, and one of D_1 , G_1 , L_1 and M_1 is NR_a , one of D_1 , G_1 , L_1 and M_1 is CR_a

and the remainder are independently selected from the group

consisting of CR_a and N, wherein R_a is as defined above;

b is 1 and D_2 , G_2 , J_2 , L_2 and M_2 are each independently selected from the

-107-

group


consisting of CR_a and N, provided that at least two of D_2 , G_2 , J_2 , L_2 and M_2 are CR_a ; or

b is 0, and one of D_2 , G_2 , L_2 and M_2 is NR_a , one of D_2 , G_2 , L_2 and M_2 is CR_a

and the remainder are independently selected from the group consisting of CR_a and N, wherein R_a is as defined above; and

n for each occurrence is independently an integer from 0 to 6.


2. The compound of Claim 1 wherein R_3 is H; R_1 for each occurrence is independently selected from the group consisting of F, Cl, Br, I, CH_3 , NO_2 , OCF_3 , OCH_3 , CN, CO_2CH_3 , CF_3 , $-CH_2NR_dR_e$, t-butyl, pyridyl, substituted or unsubstituted oxazolyl, substituted or unsubstituted benzyl, substituted or unsubstituted benzenesulfonyl, substituted or unsubstituted phenoxy, substituted or unsubstituted phenyl, substituted or unsubstituted amino, carboxyl, substituted or unsubstituted tetrazolyl, and substituted or unsubstituted styryl.
3. The compound of Claim 1 wherein R_3 is H; R_a for each occurrence is independently selected from the group consisting of F, Cl, Br, I, CH_3 , NO_2 , OCF_3 , OCH_3 , CN, CO_2CH_3 , CF_3 , t-butyl, pyridyl, substituted or unsubstituted oxazolyl, substituted or unsubstituted benzyl, substituted or unsubstituted benzenesulfonyl, substituted or unsubstituted phenoxy, substituted or unsubstituted phenyl, substituted or unsubstituted amino, carboxyl, substituted or unsubstituted tetrazolyl, and substituted or unsubstituted styryl.
4. The compound of Claim 1 wherein R_3 is H; R_2 is of the formula



wherein n is 1, 2 or 3.

-108-


5. The compound of Claim 1 wherein R_3 is H; R_2 is of the formula



wherein m is 0, 1, 2 or 3 and

R_9 is H or $-(CH_2)_pN(R_4)R_5$, wherein p is an integer from 2 to 6 and R_4 and R_5 are each, independently, H, azabicycloalkyl or Y-Z, wherein Y is selected from the group consisting of $-C(O)-$, $-(CH_2)_q-$, $-S(O)_2-$, $-C(O)O-$, $-SO_2NH-$, $-CONH-$, $-(CH_2)_qO-$, $-(CH_2)_qNH-$, and $-(CH_2)_qS(O)_r-$; wherein q is an integer from 0 to 6; and r is 0, 1 or 2; and Z is a substituted or unsubstituted moiety selected from the group consisting of alkyl, alkoxy, amino, aryl, heteroaryl and heterocycloalkyl group or R_4 , R_5 and the nitrogen atom to which they are attached together form a 3, 4, 5, 6 or 7-membered, substituted or unsubstituted heterocyclic or heterobicyclic group.

6. The compound of Claim 1 wherein R_3 is H; R_2 is of the formula



wherein m is 0, 1, 2 or 3

a and b are each, independently, an integer from 0 to 6;

Q is $-OR_6$ or $-NR_4R_5$;

each R_4 and R_5 is, independently, H, azabicycloalkyl or Y-Z, wherein Y is selected from the group consisting of $-C(O)-$, $-(CH_2)_q-$, $-S(O)_2-$, $-C(O)O-$, $-SO_2NH-$, $-CONH-$, $(CH_2)_qO-$, $-(CH_2)_qNH-$, and $-(CH_2)_qS(O)_r-$; wherein q is an integer from 0 to 6; and r is 0, 1 or 2; and Z is a substituted or unsubstituted alkyl, substituted or unsubstituted alkoxy, amino, aryl,

-109-


heteroaryl or heterocycloalkyl group or

R_4 , R_5 and the nitrogen atom to which they are attached together form a 3, 4, 5, 6 or 7-membered, substituted or unsubstituted heterocyclic or heterobicyclic group; and

5 R_6 is hydrogen or a substituted or unsubstituted alkyl group.

7. The compound of Claim 1 wherein R_3 is H; R_2 is of the formula

10



wherein n is 1, 2 or 3; and

15

R_4 is H, azabicycloalkyl or Y-Z, wherein Y is selected


from the group consisting of $-C(O)-$, $-(CH_2)_q-$, $-S(O)_2-$, $-C(O)O-$, $-SO_2NH-$, $-CONH-$, $(CH_2)_qO-$, $-(CH_2)_qNH-$, and $-(CH_2)_qS(O)_r-$;

wherein q is an integer 0 to 6; and r is 0, 1 or 2; and Z is a substituted or unsubstituted alkyl, substituted or unsubstituted amino, aryl, substituted or unsubstituted heteroaryl or substituted or unsubstituted heterocycloalkyl group.

20

8. The compound of Claim 1 wherein R_3 is H; R_2 is of the formula

25



30

wherein

m is 0, 1, 2 or 3;

R_5 is H, azabicycloalkyl or Y-Z, wherein Y is selected from the group


-110-

consisting of a covalent bond, $-C(O)-$, $-(CH_2)_q-$, $-S(O)_2-$, $-C(O)O-$, $-SO_2NH-$, $-CONH-$, $-(CH_2)_qO-$, $-(CH_2)_qNH-$, $-(CH_2)_qC(O)-$, $-C(O)(CH_2)_q-$ and $-(CH_2)_qS(O)_r-$, where the alkyl portion of $-(CH_2)_q-$, $-(CH_2)_qO-$, $-(CH_2)_qNH-$, $-(CH_2)_qC(O)-$, $-C(O)(CH_2)_q-$ and $-(CH_2)_qS(O)_r$ is optionally substituted by a halogen, hydroxy or an alkyl group; wherein q is an integer from 0 to 6; and r is 0, 1 or 2; and Z is a substituted or unsubstituted alkyl, substituted or unsubstituted amino, substituted or unsubstituted alkoxy, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl or substituted or unsubstituted heterocycloalkyl group;

or Y and Z together are a natural or unnatural amino acid, which may be mono- or di-alkylated at the amine nitrogen; and

R_6 represents one or more substituents each independently selected from the group consisting of hydrogen, hydroxy, oxo, substituted or unsubstituted alkyl, substituted or unsubstituted aryl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted alkoxycarbonyl, substituted or unsubstituted alkoxyalkyl, substituted or unsubstituted aminocarbonyl, substituted or unsubstituted alkylcarbonyl, substituted or unsubstituted arylcarbonyl, substituted or unsubstituted heterocyclylcarbonyl, substituted or unsubstituted aminoalkyl and substituted or unsubstituted arylalkyl; provided that the carbon atoms adjacent to the nitrogen atom are not substituted by a hydroxy group.


9. The compound of Claim 1 wherein R_3 is H; R_2 is of the formula



-111-

wherein R_4 is H, substituted or unsubstituted alkyl, substituted or unsubstituted azabicycloalkyl or Y-Z, wherein Y is selected from the group consisting of -C(O)-, $-(CH_2)_q-$, $-S(O)_2-$, -C(O)O-, $-SO_2NH-$, -CONH-, $-(CH_2)_qO-$, $-(CH_2)_qNH-$, and $-(CH_2)_qS(O)_r-$; wherein q is an integer from 0 to 6, and r is 0, 1 or 2; and Z is hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted amino, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl or substituted or unsubstituted heterocycloalkyl.

10 10. The compound of Claim 1 wherein R_3 is H; R_2 is of the formula



wherein


m is an integer from 1 to 6; and

R_4 and R_5 are each, independently, H, substituted or unsubstituted azabicycloalkyl or Y-Z, wherein Y is selected from the group consisting of -C(O)-, $-(CH_2)_q-$, $-S(O)_2-$, -C(O)O-, $-SO_2NH-$, -CONH-, $-(CH_2)_qO-$, $-(CH_2)_qNH-$, and $-(CH_2)_qS(O)_r-$; wherein q is an integer from 0 to 6; and r is 0, 1 or 2; and Z is a substituted or unsubstituted alkyl, substituted or unsubstituted amino, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl or substituted or unsubstituted heterocycloalkyl group; or

R_4 , R_5 and the nitrogen atom to which they are attached together form a 3, 4, 5, 6 or 7-membered, substituted or unsubstituted heterocyclic or substituted or unsubstituted heterobicyclic group.

-112-

11. The compound of Claim 1 wherein R_3 is H; R_2 is of the formula



wherein

n is an integer from 0 to 4;

r is 0 and m is an integer from 1 to 6; or

r is 1 and m is an integer from 0 to 6;

Q is $-OR_6$ or $-NR_4R_5$;

each R_4 and R_5 is, independently, H, substituted or unsubstituted azabicycloalkyl or $Y-Z$, wherein Y is selected from the group consisting of $-C(O)-$, $-(CH_2)_q-$,


$-S(O)_2-$, $-C(O)O-$, $-SO_2NH-$, $-CONH-$, $-(CH_2)_qO-$, $-(CH_2)_qNH-$, and $-(CH_2)_qS(O)_r-$;

q is an integer from 0 to 6; and r is 0, 1 or 2; and Z is a substituted or unsubstituted alkyl, substituted or unsubstituted alkoxy, substituted or unsubstituted amino, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl or substituted or unsubstituted heterocycloalkyl group; or

R_4 , R_5 and the nitrogen atom to which they are attached together form a 3, 4, 5 or 6-membered, substituted or unsubstituted heterocyclic group; and R_6 is hydrogen or a substituted or unsubstituted alkyl group.

-113-

12. The compound of Claim 1 wherein R_3 is H; R_2 is of the formula




n is an integer from 0 to 4;

m is an integer from 0 to 6;

R_4 is H, substituted or unsubstituted azabicycloalkyl or Y-Z, wherein Y is selected from the group consisting of -C(O)-, $-(CH_2)_q-$, $-S(O)_2-$, $-C(O)O-$, $-SO_2NH-$, $-CONH-$, $-(CH_2)_qO-$, $-(CH_2)_qNH-$, and $-(CH_2)_qS(O)_r-$; wherein q is an integer from 0 to 6; and r is 0, 1 or 2; and Z is substituted or unsubstituted alkyl, substituted or unsubstituted amino, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl or substituted or unsubstituted heterocycloalkyl; and

R_6 is hydrogen or a substituted or unsubstituted alkyl group.

13. The compound of Claim 10 wherein R_4 , R_5 and the nitrogen atom together form a heterocyclic group of the formula




wherein

R_7 , R_8 , R_9 , R_{10} , R_{11} , R_{12} , R_{13} and R_{14} are each, independently, lower alkyl or hydrogen; or at least one pair of substituents R_7 and R_8 ; R_9 and R_{10} ; R_{11} and R_{12} ; or R_{13} and R_{14} together are an oxygen atom; or at least one of R_7 and R_9 is cyano, $CONHR_{15}$, $COOR_{15}$, CH_2OR_{15} or $CH_2NR_{15}(R_{16})$, wherein R_{15} and R_{16} are each, independently, H, azabicycloalkyl or V-L, wherein V is

-114-

- selected from the group consisting of $-\text{C}(\text{O})-$, $-(\text{CH}_2)_p-$, $-\text{S}(\text{O})_2-$, $-\text{C}(\text{O})\text{O}-$, $-\text{SO}_2\text{NH}-$, $-\text{CONH}-$, $(\text{CH}_2)_q\text{O}-$, $-(\text{CH}_2)_q\text{NH}-$, and $-(\text{CH}_2)_q\text{S}(\text{O})_r-$; wherein p is an integer from 0 to 6, q is an integer from 0 to 6, and r is 0, 1 or 2; and L is substituted or unsubstituted alkyl, substituted or unsubstituted amino, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl or substituted or unsubstituted heterocycloalkyl; or R_{15} , R_{16} and the nitrogen atom together form a 3, 4, 5, 6 or 7-membered, substituted or unsubstituted heterocyclic or a substituted or unsubstituted heterobicyclic group;
- X is O , S , SO , SO_2 , CH_2 , CHOR_{17} or NR_{17} , wherein R_{17} is hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted aryl, substituted or unsubstituted arylalkyl, $-\text{C}(\text{NH})\text{NH}_2$, $-\text{C}(\text{O})\text{R}_{17}$, or $-\text{C}(\text{O})\text{OR}_{18}$, wherein R_{18} is hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted aryl or substituted or unsubstituted arylalkyl; and
- t is 0 or 1.
14. The compound of Claim 10 wherein R_4 , R_5 and the nitrogen atom together form a heterocycle of the formula



wherein

- R_{19} and R_{20} are each, independently, hydrogen or lower alkyl; or R_{19} and R_{20} together are an oxygen atom;
- R_{21} and R_{22} are each, independently, H, substituted or unsubstituted azabicycloalkyl or V-L, wherein V is selected from the group consisting of $-\text{C}(\text{O})-$, $-(\text{CH}_2)_p-$, $-\text{S}(\text{O})_2-$, $-\text{C}(\text{O})\text{O}-$, $-\text{SO}_2\text{NH}-$, $-\text{CONH}-$, $(\text{CH}_2)_q\text{O}-$, $(\text{CH}_2)_q\text{NH}-$, and $-(\text{CH}_2)_q\text{S}(\text{O})_r-$; wherein p is an integer from 0 to 6, q is an

-115-


integer from 0 to 6, and r is 0, 1 or 2; and L is substituted or unsubstituted alkyl, substituted or unsubstituted amino, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl or substituted or unsubstituted heterocycloalkyl group; or

- 5 R_{21} , R_{22} and the nitrogen atom together form a 3, 4, 5 or 6-membered, substituted or unsubstituted heterocyclic group;

m is an integer from 1 to 6; and

n is an integer from 0 to 6.

- 10 15. The compound of Claim 10 wherein R_4 , R_5 and the nitrogen atom together form a heterocyclic group of the formula



15


wherein

m is an integer from 1 to 6; and

R_{23} is CH_2OH , NRR' , $\text{C(O)NRR}'$ or COOR , wherein R and R' are each, independently, hydrogen or substituted or unsubstituted alkyl, substituted or unsubstituted aryl or substituted or unsubstituted arylalkyl.

20

16. The compound of Claim 10 wherein R_4 , R_5 and the nitrogen atom together form a heterocyclic group of the formula



25


wherein R_{24} is substituted or unsubstituted alkyl, substituted or unsubstituted aryl or substituted or unsubstituted arylalkyl, carboxyl, cyano, C(O)OR_{25} , $\text{CH}_2\text{OR}_{25}$, $\text{CH}_2\text{NR}_{26}\text{R}_{27}$ or C(O)NHR_{26} , wherein R_{25} is substituted or unsubstituted alkyl, substituted or unsubstituted aryl, substituted or unsubstituted arylalkyl, substituted or unsubstituted heterocyclic or

30

-116-

substituted or unsubstituted heterocycloaryl; and R_{26} and R_{27} are each, independently, H, substituted or unsubstituted azabicycloalkyl or V-L, wherein V is selected from the group consisting of $-C(O)-$, $-(CH_2)_p-$, $-S(O)_2-$, $-C(O)O-$, $-SO_2NH-$, $-CONH-$, $(CH_2)_qO-$, $-(CH_2)_qNH-$, and $-(CH_2)_qS(O)_r-$; wherein p is an integer from 0 to 6, q is an integer from 0 to 6, and r is 0, 1 or 2; and L is substituted or unsubstituted alkyl, substituted or unsubstituted amino, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl or substituted or unsubstituted heterocycloalkyl; or R_{26} , R_{27} and the nitrogen atom together form a 3, 4, 5 or 6-membered, substituted or unsubstituted heterocyclic group.

17. The compound of Claim 10 wherein at least one of R_4 and R_5 is of the formula Y-Z, wherein Z is of the formula



wherein


T is C(O), S, SO, SO_2 , CHOR or NR, wherein R is hydrogen or a substituted or unsubstituted alkyl, substituted or unsubstituted aryl or substituted or unsubstituted arylalkyl group; and n is 0, 1 or 2.

18. The compound of Claim 10 wherein at least one of R_4 and R_5 is of the formula Y-Z, wherein Z is of the formula $-N(R_{28})R_{29}$, wherein R_{28} and R_{29} are each, independently, substituted or unsubstituted carboxyalkyl, substituted or unsubstituted alkoxycarbonylalkyl, substituted or unsubstituted hydroxyalkyl, substituted or unsubstituted alkylsulfonyl, substituted or unsubstituted alkylcarbonyl or substituted or unsubstituted cyanoalkyl; or R_{28} and R_{29} , together with the nitrogen atom, form a five- or six-membered

-117-

substituted or unsubstituted heterocyclic group.

19. The compound of Claim 11 wherein R_4 , R_5 and the nitrogen atom together form a heterocycle of the formula




10 wherein

- R_7 , R_8 , R_9 , R_{10} , R_{11} , R_{12} , R_{13} and R_{14} are each, independently, lower alkyl or hydrogen; or at least one pair of substituents R_7 and R_8 ; R_9 and R_{10} ; R_{11} and R_{12} ; or R_{13} and R_{14} together are an oxygen atom; or at least one of R_7 and R_9 is cyano, CONHR_{15} , COOR_{15} , $\text{CH}_2\text{OR}_{15}$ or $\text{CH}_2\text{NR}_{15}(\text{R}_{16})$, wherein R_{15} and R_{16} are each, independently, H, substituted or unsubstituted azabicycloalkyl or V-L, wherein V is selected from the group consisting of $-\text{C}(\text{O})-$, $-(\text{CH}_2)_p-$, $-\text{S}(\text{O})_2-$, $-\text{C}(\text{O})\text{O}-$, $-\text{SO}_2\text{NH}-$, $-\text{CONH}-$, $(\text{CH}_2)_q\text{O}-$, $-(\text{CH}_2)_q\text{NH}-$, and $-(\text{CH}_2)_q\text{S}(\text{O})_r-$; wherein p is an integer from 0 to 6, q is an integer from 0 to 6, and r is 0, 1 or 2; and L is substituted or unsubstituted alkyl, substituted or unsubstituted amino, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl or substituted or unsubstituted heterocycloalkyl; or R_{15} , R_{16} and the nitrogen atom together form a 3, 4, 5, 6 or 7-membered, substituted or unsubstituted heterocyclic or heterobicyclic group;
- 25 X is O, S, SO, SO_2 , CH_2 , CHOR_{17} or NR_{17} , wherein R_{17} is hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted aryl, substituted or unsubstituted arylalkyl, $-\text{C}(\text{NH})\text{NH}_2$, $-\text{C}(\text{O})\text{R}_{18}$, or $-\text{C}(\text{O})\text{OR}_{18}$, wherein R_{18} is hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted aryl or substituted or unsubstituted arylalkyl; and
- 30 t is 0 or 1.

-118-

20. The compound of Claim 11 wherein R_4 , R_5 and the nitrogen atom together form a heterocycle of the formula

5



wherein

10

R_{19} and R_{20} are each, independently, hydrogen or lower alkyl; or R_{19} and R_{20} together are an oxygen atom;

15

R_{21} and R_{22} are each, independently, H, substituted or unsubstituted azabicycloalkyl or V-L, wherein V is selected from the group consisting of -C(O)-, $-(CH_2)_p-$, $-S(O)_2-$, $-C(O)O-$, $-SO_2NH-$, $-CONH-$, $(CH_2)_qO-$, $-(CH_2)_qNH-$, and $-(CH_2)_qS(O)_r-$; wherein p is an integer from 0 to 6, q is an integer from 0 to 6, and r is 0, 1 or 2; and L is substituted or unsubstituted alkyl, substituted or unsubstituted amino, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl or substituted or unsubstituted heterocycloalkyl group; or

20

R_{21} , R_{22} and the nitrogen atom together form a 3, 4, 5 or 6-membered, substituted or unsubstituted heterocyclic group;


m is an integer from 1 to 6; and

n is an integer from 0 to 6.

25 21.

The compound of Claim 11 wherein R_4 , R_5 and the nitrogen atom together form a heterocyclic group of the formula

30




-119-

wherein

m is an integer from 1 to 6; and

R₂₃ is CH₂OH, NRR', C(O)NRR' or COOR, wherein R is hydrogen or a substituted or unsubstituted alkyl, substituted or unsubstituted aryl or substituted or unsubstituted arylalkyl group.


22. The compound of Claim 11 wherein R₄, R₅ and the nitrogen atom together form a heterocyclic group of the formula



wherein R₂₄ is substituted or unsubstituted alkyl, substituted or unsubstituted aryl or substituted or unsubstituted arylalkyl, carboxyl, cyano, C(O)OR₂₅, CH₂OR₂₅, CH₂NR₂₆R₂₇ or C(O)NHR₂₆, wherein R₂₅ is substituted or unsubstituted alkyl, substituted or unsubstituted aryl, substituted or unsubstituted arylalkyl, substituted or unsubstituted heterocyclic or substituted or unsubstituted heterocycloaryl group; and R₂₆ and R₂₇ are each, independently, H, substituted or unsubstituted azabicycloalkyl or V-L, wherein V is selected from the group consisting of -C(O)-, -(CH₂)_p-, -S(O)₂-, -C(O)O-, -SO₂NH-, -CONH-, (CH₂)_qO-, -(CH₂)_qNH-, and -(CH₂)_qS(O)_r-; wherein p is an integer from 0 to 6, q is an integer from 0 to 6, and r is 0, 1 or 2; and L is substituted or unsubstituted alkyl, substituted or unsubstituted amino, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl or substituted or unsubstituted heterocycloalkyl group; or R₂₆, R₂₇ and the nitrogen atom together form a 3, 4, 5 or 6-membered, substituted or unsubstituted heterocyclic group.

23. The compound of Claim 11 wherein at least one of R₄ and R₅ is of the formula Y-Z, wherein Z is of the formula

-120-



5

wherein

g is 0 or 1;

T is $C(O)$, O , S , SO , SO_2 , CH_2 , $CHOR_{17}$ or NR_{17} , wherein R_{17} is hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted aryl,

10

substituted or unsubstituted arylalkyl, $-C(NH)NH_2$, $-C(O)R_{18}$, or $-C(O)OR_{18}$,

wherein R_{18} is hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted aryl or substituted or unsubstituted arylalkyl; and

R_{32} is hydrogen, cyano, substituted or unsubstituted alkyl, substituted or unsubstituted alkoxy carbonyl, substituted or unsubstituted alkoxyalkyl,

15

substituted or unsubstituted hydroxyalkyl, substituted or unsubstituted aminocarbonyl, substituted or unsubstituted alkylcarbonyl or substituted or unsubstituted arylalkyl.

24. The compound of Claim 11 wherein at least one of R_4 and R_5 is of the

20

formula $Y-Z$, wherein Z is of the formula $-N(R_{28})R_{29}$, wherein R_{28} and R_{29} are each, independently, substituted or unsubstituted carboxyalkyl, substituted or unsubstituted alkoxy carbonylalkyl, substituted or unsubstituted

hydroxyalkyl, substituted or unsubstituted alkylsulfonyl, substituted or unsubstituted alkylcarbonyl or substituted or unsubstituted cyanoalkyl; or

25

R_{28} and R_{29} , together with the nitrogen atom, form a five- or six-membered substituted or unsubstituted heterocyclic group.

25. The compound of Claim 8 wherein R_5 is $Y-Z$, wherein Z is of the formula

30

$N(R_{30})R_{31}$, wherein R_{30} and R_{31} are each, independently, hydrogen, alkyl, alkoxy carbonyl, alkoxyalkyl, hydroxyalkyl, aminocarbonyl, cyano, alkylcarbonyl or arylalkyl.

-121-

26. The compound of Claim 8 wherein R_5 is Y-Z, wherein Z is of the formula



wherein

- 10 each X is, independently, CH or N; and
 R_{32} is hydrogen, cyano, substituted or unsubstituted alkyl, substituted or unsubstituted alkoxy carbonyl, substituted or unsubstituted alkoxy alkyl, substituted or unsubstituted hydroxy alkyl, substituted or unsubstituted aminocarbonyl, substituted or unsubstituted alkyl carbonyl or substituted or unsubstituted aryl alkyl group.
- 15

27. The compound of Claim 8 wherein R_5 is Y-Z, wherein Z is of the formula




wherein

- g is 0 or 1;
- 25 T is O, S, SO, SO₂, CH₂, CHOR₁₇ or NR₁₇, wherein R₁₇ is hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted aryl, substituted or unsubstituted aryl alkyl, C(O)NH₂, -C(NH)NH₂, -C(O)R₁₇, or -C(O)OR₁₈, wherein R₁₈ is hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted aryl or substituted or unsubstituted aryl alkyl; and
- 30 R_{32} is hydrogen, cyano, substituted or unsubstituted alkyl, substituted or unsubstituted alkoxy carbonyl, substituted or unsubstituted alkoxy alkyl,

-122-

substituted or unsubstituted hydroxyalkyl, substituted or unsubstituted aminocarbonyl, substituted or unsubstituted alkylcarbonyl or substituted or unsubstituted arylalkyl group.

- 5 28. The compound of Claim 8 wherein R_5 is Y-Z, wherein Z is of the formula




10 wherein

g is 0, 1 or 2; and

R_{32} is hydrogen, cyano, substituted or unsubstituted alkyl, substituted or unsubstituted alkoxycarbonyl, substituted or unsubstituted alkoxyalkyl, substituted or unsubstituted hydroxyalkyl, substituted or unsubstituted

15 aminocarbonyl, substituted or unsubstituted alkylcarbonyl or substituted or unsubstituted arylalkyl group.

29. The compound of Claim 8 wherein R_5 is Y-Z, wherein Z is of the formula



20

wherein


T is C(O), O, S, SO, SO₂, CH₂, CHOR₁₇ or NR₁₇, wherein R₁₇ is hydrogen, substituted or unsubstituted alkyl, aryl, arylalkyl, -C(NH)NH₂, -C(O)R₁₈, or -C(O)OR₁₈, wherein R₁₈ is hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted aryl or substituted or unsubstituted arylalkyl; g is 0 or 1; and

25 30 R_{32} is hydrogen, cyano, substituted or unsubstituted alkyl, substituted or unsubstituted alkoxycarbonyl, substituted or unsubstituted alkoxyalkyl, substituted or unsubstituted hydroxyalkyl, substituted or unsubstituted

-123-

aminocarbonyl, substituted or unsubstituted alkylcarbonyl or substituted or unsubstituted arylalkyl group.


30. The compound of Claim 8 wherein R_5 is Y-Z, wherein Z is of the formula



10 wherein

R_{32} is hydrogen, cyano, substituted or unsubstituted alkyl, substituted or unsubstituted alkoxycarbonyl, substituted or unsubstituted alkoxyalkyl, substituted or unsubstituted hydroxyalkyl, substituted or unsubstituted aminocarbonyl, alkylcarbonyl, substituted or unsubstituted thioalkoxy or substituted or unsubstituted arylalkyl; and R_{33} is hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted alkoxycarbonyl, substituted or unsubstituted alkoxyalkyl, substituted or unsubstituted aminocarbonyl, perhaloalkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkylcarbonyl or substituted or unsubstituted arylalkyl.

31. The compound of Claim 1 wherein R_3 is H; R_2 is of the formula



wherein

m is 0 or 1;


R_{34} , R_{35} , R_{36} , R_{37} , R_{38} , R_{39} , R_{40} and R_{41} are each, independently, methyl or

-124-

hydrogen; or at least one pair of substituents R_{34} and R_{35} ; R_{36} and R_{37} ; R_{38} and R_{39} ; or R_{40} and R_{41} together are an oxygen atom; and

R_{42} is H, substituted or unsubstituted azabicycloalkyl or Y-Z, wherein Y is selected from the group consisting of -C(O)-, $-(CH_2)_p-$, $-S(O)_2-$, -C(O)O-, -SO₂NH-, -CONH-, $(CH_2)_qO-$, $-(CH_2)_qNH-$, and $-(CH_2)_qS(O)_r-$; wherein p is an integer from 0 to 6, q is an integer from 0 to 6, and r is 0, 1 or 2; and Z is substituted or unsubstituted alkyl, substituted or unsubstituted amino, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl or substituted or unsubstituted heterocycloalkyl group; or

R_{42} is of the formula



wherein

u is 0 or 1;

R_{43} , R_{44} , R_{45} , R_{46} , R_{47} , R_{48} , R_{49} and R_{50} are each, independently, methyl or hydrogen;


or at least one pair of substituents R_{43} and R_{44} ; R_{45} and R_{46} ; R_{47} and R_{48} ; or R_{49} and R_{50} together are an oxygen atom; and

R_{51} is H, substituted or unsubstituted azabicycloalkyl or V-L, wherein V is selected from the group consisting of -C(O)-, $-(CH_2)_p-$, $-S(O)_2-$, -C(O)O-, -SO₂NH-, -CONH-, $(CH_2)_qO-$, $-(CH_2)_qNH-$, and $-(CH_2)_qS(O)_r-$; wherein p is an integer from 0 to 6, q is an integer from 0 to 6, and r is 0, 1 or 2; and L is substituted or unsubstituted alkyl, substituted or unsubstituted amino, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl or substituted or unsubstituted heterocycloalkyl.

-125-

32. The compound of Claim 1 wherein R_3 is H; R_2 is of the formula

5



wherein

10

h, i, j, k and l are independently 0 or 1;

$R_{52}, R_{53}, R_{54}, R_{55}, R_{56}, R_{57}, R_{58}, R_{59}, R_g$ and R_h are each, independently, methyl or hydrogen; or at least one pair of substituents R_{52} and R_{53} ; R_{54} and R_{55} ; R_{56} and R_{57} ; or R_{58} and R_{59} together are an oxygen atom; and


15

R_{60} is H, substituted or unsubstituted azabicycloalkyl or Y-Z, wherein Y is selected from the group consisting of $-C(O)-$, $-(CH_2)_p-$, $-S(O)_2-$, $-C(O)O-$, $-SO_2NH-$, $-CONH-$, $(CH_2)_qO-$, $-(CH_2)_qNH-$, and $-(CH_2)_qS(O)_r-$; wherein p is an integer from 0 to 6, q is an integer from 0 to 6, and r is 0, 1 or 2; and Z is substituted or unsubstituted alkyl, substituted or unsubstituted amino, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl or substituted or unsubstituted heterocycloalkyl; or

20

R_{60} is of the formula

25



wherein

v is 0 or 1;

30

$R_{61}, R_{62}, R_{63}, R_{64}, R_{65}, R_{66}, R_{67}$ and R_{68} are each, independently, lower alkyl or hydrogen; or at least one pair of substituents R_{61} and R_{62} ; R_{63} and R_{64} ; R_{65} and R_{66} ; and R_{67} and R_{68} together are an oxygen atom; and

-126-

- 5 R_{69} is H, substituted or unsubstituted azabicycloalkyl or V-1, wherein V is selected from the group consisting of -C(O)-, -(CH₂)_p-, -S(O)₂-, -C(O)O-, -SO₂NH-, -CONH-, (CH₂)_qO-, -(CH₂)_qNH-, and -(CH₂)_qS(O)_r-; wherein p is an integer from 0 to 6, q is an integer from 0 to 6, and r is 0, 1 or 2; and L is substituted or unsubstituted alkyl, substituted or unsubstituted amino, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl or substituted or unsubstituted heterocycloalkyl.
- 10 33. A method of inhibiting one or more protein kinase activity in a patient comprising administering a therapeutically effective amount of a compound of Claim 1 or a physiologically acceptable salt, prodrug or biologically active metabolites thereof to said patient.
- 15 34. The method of Claim 33 wherein said protein kinase is selected from the group consisting of KDR, FGFR-1, PDGFR β , PDGFR α , IGF-1R, c-Met, Flt-1, Flt-4, TIE-2, TIE-1, Lck, Src, fyn, Lyn, Blk, hck, fgr and yes.
- 20 35. A method of affecting hyperproliferative disorders in a patient comprising administering a therapeutically effective amount of a compound of Claim 1 or a physiologically acceptable salt, prodrug or biologically active metabolites thereof to said patient.
- 25 36. A method of affecting angiogenesis in a patient comprising administering a therapeutically effective amount of a compound of Claim 1 or a physiologically acceptable salt, prodrug or biologically active metabolites thereof to said patient.
- 30 37. The method of Claim 33 wherein the protein kinase is a protein serine/threonine kinase or a protein tyrosine kinase.

-127-

38. A method of treating one or more ulcers in a patient comprising administering a therapeutically effective amount of a compound of Claim 1 or a physiologically acceptable salt, prodrug or biologically active metabolites thereof to said patient.
- 5
39. The method of Claim 38 wherein the ulcer or ulcers are caused by a bacterial or fungal infection; or the ulcer or ulcers are Mooren ulcers; or the ulcer or ulcers are a symptom of ulcerative colitis.
- 10
40. A method of treating a condition in a patient comprising administering a therapeutically effective amount of a compound of Claim 1 or a physiologically acceptable salt, prodrug or biologically active metabolites thereof to said patient, wherein said condition is an ocular condition, a cardiovascular condition, a cancer, Crow-Fukase (POEMS) syndrome, a
- 15
- diabetic condition, sickle cell anaemia, chronic inflammation, systemic lupus, glomerulonephritis, synovitis, inflammatory bowel disease, Crohn's disease, glomerulonephritis, rheumatoid arthritis, osteoarthritis, multiple sclerosis, graft rejection, Lyme disease, sepsis, von Hippel Lindau disease, pemphigoid, psoriasis, Paget's disease, polycystic kidney disease, fibrosis,
- 20
- sarcoidosis, cirrhosis, thyroiditis, hyperviscosity syndrome, Osler-Weber-Rendu disease, chronic occlusive pulmonary disease, asthma or edema following burns, trauma, radiation, stroke, hypoxia, ischemia, ovarian hyperstimulation syndrome, preeclampsia, menometrorrhagia, endometriosis, or infection by Herpes simplex, Herpes Zoster, human immunodeficiency
- 25
- virus, parapoxvirus, protozoa or toxoplasmosis.
41. The method of Claim 40 wherein the ocular condition is ocular or macular edema, ocular neovascular disease, scleritis, radial keratotomy, uveitis, vitritis, myopia, optic pits, chronic retinal detachment, post-laser treatment
- 30
- complications, conjunctivitis, Stargardt's disease, Eales disease, retinopathy or macular degeneration.

-128-


42. The method of Claim 40 wherein the cardiovascular condition is atherosclerosis, restenosis, ischemia/reperfusion injury, vascular occlusion or carotid obstructive disease.
- 5
43. The method of Claim 40 wherein the cancer is a solid tumor, a sarcoma, fibrosarcoma, osteoma, melanoma, retinoblastoma, a rhabdomyosarcoma, glioblastoma, neuroblastoma, teratocarcinoma, an hematopoietic malignancy, Kaposi's sarcoma, Hodgkin's disease, lymphoma, myeloma, leukemia or
- 10 malignant ascites.
44. The method of Claim 40 wherein the diabetic condition is insulin-dependent diabetes mellitus glaucoma, diabetic retinopathy or microangiopathy.
- 15 45. A method of decreasing fertility in a patient, said method comprising the step of administering to the patient an effective amount of a compound of Claim 1 or a physiologically acceptable salt, prodrug or biologically active metabolite thereof.
- 20 46. The method of Claim 36 wherein the compound or a physiologically acceptable salt, prodrug or biologically active metabolite thereof is administered in an amount effective to promote angiogenesis or vasculogenesis.
- 25 47. The method of Claim 34 wherein the protein kinase is Tie-2.
48. The method of Claim 46 wherein the compound of Formula I, or physiologically acceptable salt, prodrug or biologically active metabolite thereof, is administered in combination with a pro-angiogenic growth factor.
- 30

-129-

49. The method of Claim 48 wherein the pro-angiogenic growth factor is selected from the group consisting of VEGF, VEGF-B, VEGF-C, VEGF-D, VEGF-E, HGF, FGF-1, FGF-2, derivatives thereof and anti-idiotypic antibodies.
50. The method of Claim 46 wherein the patient is suffering from anemia, ischemia, infarct, transplant rejection, a wound, gangrene or necrosis.
51. The method of Claim 33 wherein the protein kinase activity is involved in T cell activation, B cell activation, mast cell degranulation, monocyte activation, the potentiation of an inflammatory response or a combination thereof.
52. A compound according to Claim 1, wherein R_3 is H;
 R_2 is $-Z^{101}-Z^{102}$ where Z^{101} is a covalent bond, $-(C_1-C_6)-$, $-(C_1-C_6)-O-$, $-(C_1-C_6)-C(O)-$, $-(C_1-C_6)-C(O)O-$, $-(C_1-C_6)-C(O)-NH-$, $-(C_1-C_6)-C(O)-N((C_1-C_6))-$ or a substituted phenyl group; and
 Z^{102} is hydrogen, a substituted or unsubstituted alkyl group or a substituted or unsubstituted, saturated or unsaturated heterocyclic group.
53. A compound according to Claim 52, wherein Z^{101} is selected from the group consisting of $-CH_2-C(O)O-$, $-CH_2-C(O)-$, $-CH_2-C(O)-NH-$, $-CH_2-C(O)-N(Me)-$, $-CH(Me)-C(O)O-$, $-(CH_2)_3-C(O)O-$, $-CH(Me)-C(O)-NH-$, and $-(CH_2)_3-C(O)-NH-$;
 Z^{102} is selected from the group consisting of hydrogen, methyl, ethyl, N,N-dimethylaminoethyl, N,N-diethylaminoethyl, 2-phenyl-2-hydroxyethyl, morpholino, piperazinyl, N-methylpiperazinyl and 2-hydroxymethylpyrrolidinyl.

-130-


54. A compound according to Claim 53, wherein R_1 is



Z^{100} is a substituted or unsubstituted benzoxazolyl or a substituted or unsubstituted benzthiazolyl.

5

55. A compound according to Claim 8, 9, 10 or 53, wherein R_1 is




10 where there is only one R_a and it is H or F.


56. A compound according to Claim 52, wherein Z^{101} is a covalent bond; and Z^{102} is an optionally substituted pyridyl.

15 57. A compound according to Claim 56, wherein R_1 is

-131-




58. A compound according to Claim 1, wherein R_3 is H;
 R_2 is cyclopentyl; and



59. A compound according to Claim 58, wherein
 Z^{110} is hydrogen;
A is O; and Z^{100} is optionally substituted phenyl, furanyl or thienyl, where
10 Z^{100} is optionally substituted with one or more substituents each
independently selected from the group consisting of F, COOH, NO₂, OMe, -
COOMe, OCF₃ and CF₃.
60. A compound according to Claim 58, wherein
15 Z^{110} is hydrogen;
A is -O-, -O-(CR₂)_n-C(O)- or -O-(CR₂)_n-O-;
n for each occurrence is 0 to 3;
 Z^{100} is an optionally substituted group selected from the group consisting of
cyclohexyl, phenyl, tetrahydropyranyl, tetrahydrofuranyl, isoxazolyl and
20 piperidiny; where Z^{100} is optionally substituted with one or more substituents
selected from the group consisting of alkyl, alkoxy, halo, hydroxy and
alkoxycarbonyl.
61. A compound according to Claim 58, wherein R^2 is an optionally substituted
25 group selected from the group consisting of cyclobutyl and cyclohexyl.


-132-

62. A compound according to Claim 61, wherein R^2 is optionally substituted with one or more substituents selected from the group consisting of hydroxy, alkyl, hydroxyalkyl, carboxyalkyl and phenylalkoxyalkyl.
- 5 63. A compound according to Claim 62, wherein R_1 is 4-phenoxyphenyl.
64. A compound according to Claim 6 wherein
m is 2; a is 0; R_6 is H; b is 1 or 2; and R_4 and R_5 are each hydrogen.
- 10 65. A compound according to Claim 8, wherein
m is 0, 1 or 2; R_6 is hydrogen; R_5 is H or Y-Z;
where Y is a covalent bond, $-C(O)-$, $-(CH_2)_qO-$, $-(CH_2)_q-$, $-(CH_2)_qC(O)-$ or $-C(O)(CH_2)_q-$, where the alkyl portion of $-(CH_2)_qO-$, $-(CH_2)_q-$, $-(CH_2)_qC(O)-$ and $-C(O)(CH_2)_q-$ is optionally substituted by a halogen, hydroxy or an alkyl
15 group; and
Z is hydrogen, alkyl, optionally substituted alkyl, alkoxyalkyl, optionally substituted heterocycloalkyl, optionally substituted heteroaryl, or optionally substituted amino.
- 20 66. A compound according to Claim 65, wherein
Z is hydrogen, methyl, ethyl, hydroxymethyl, methoxyethyl, N-methylpiperidinyl, (t-butoxycarbonyl)(hydroxy)-piperidinyl, hydroxypiperidinyl, (hydroxymethyl)piperidinyl, (hydroxy)(methyl)-piperidinyl, morpholino, (methoxyethyl)piperizinyl, methylpiperizinyl, 4-piperidinylpiperidinyl,
25 imidazolyl, methylimidazolyl, N-methylamino, N,N-dimethylamino, N-isopropylamino, N,N-diethylamino, 2,3-dihydroxypropylamino, 2-hydroxyethylamino, 3-hydroxypropylamino, methoxyethylamino, ethoxycarbonylmethylamino, phenylmethylamino, N-methyl-N-methoxyamino, , furanylethylamino, piperidineethylamino,
30 N-(2-N,N-dimethylaminoethyl)-N-methylamino, 2-N,N-


-133-

dimethylaminoethylamino, N-methyl-N-(N-methylpiperidin-4-yl)amino, 2-morpholino-ethylamino, 3-morpholino-propylamino, 3-imidazolylpropylamino, or 3-(2-oxopyrrolidinyl)propylamino.

- 5 67. A compound according to Claim 8, wherein m is 2; R_5 is Y-Z; Y is -C(O)-;



68. A compound according to Claim 9, wherein R_4 is hydrogen or methyl;



A is selected from the group consisting of O, -N(R)- and -N(R)C(O)-;

Z^{III} is $-(CH_2)_n$ -cycloalkyl- $(CH_2)_n$;

R is hydrogen or alkyl;

n is 0 to 5;

- 15 R_a is one or more substituents each independently selected from the group consisting of H, OH, F, Cl, methyl and methoxy; and


R_b is one or more substituents each independently selected from the group consisting of H, CN, F, CF_3 , OCF_3 , methyl, methoxy and an optionally substituted amino group;

- 20 where said amino group is optionally substituted with one or two groups each independently selected from the group consisting of alkyl, alkoxyalkyl, phenyl, substituted phenyl, and optionally substituted heteroaryl.

- 25 69. A compound according to Claim 68, wherein R_b is 4-methylphenylthio or 2-pyridinylthio.

-134-

70. A compound according to Claim 9, wherein




R_1 is

where Z^{100} is selected from the group consisting of benzo[b]thiophene, furanyl and thiophene.

5

71. A compound according to Claim 9C, wherein R_a is alkoxy; A is $-\text{NH}-\text{C}(\text{O})-$; and there is a covalent bond between A and Z^{100} .

72. A compound according to Claims 1, 8 or 9, wherein



R_1 is

A is selected from the group consisting of $-\text{N}(\text{R})-\text{C}(\text{O})-\text{N}(\text{R})-$, $-(\text{CH}_2)_n-$, $\text{N}(\text{R})\text{C}(\text{O})\text{N}(\text{R})-$, $-\text{N}(\text{R})-$ and $-\text{N}(\text{R})-\text{SO}_2-$; R is hydrogen or alkyl;

Z^{100} is , pyridinyl, thiazolyl, furanyl, benzofuranyl or oxazolyl;

15

X is S, O or NR where R for each occurrence is independently H or Me;

R_a is one or more substituents each independently selected from the group consisting of H and F; and

R_b is one or more substituents each independently selected from the group consisting of H, F, Cl, Br, NO_2 , CF_3 , alkyl, alkoxy and alkoxycarbonyl.

20


73. A compound according to Claim 72, wherein

R_4 is methyl; m is 1, 2 or 3; R_5 is Y-Z, where Y is $-\text{C}(\text{O})\text{O}-$, $-\text{C}(\text{O})-$ or $-\text{C}(\text{O})-(\text{CH}_2)_p-$; and Z is aminoalkyl, N-alkylamino, N,N-dialkylamino or

-135-


hydroxyalkylaminoalkyl.

74. A compound according to Claim 9, wherein

 R_4 is methyl; R_1 is

where n is 0 to 3; Z^{100} is an optionally substituted group selected from the group consisting of indolyl, indenyl, methylindenyl, methylindolyl, dimethylaminophenyl, phenyl, cyclohexyl and benzofuranyl.

75. A compound according to claim 9, wherein



Z^{100} is an optionally substituted group selected from the group consisting of phenyl, imidazolyl, indolyl, furanyl, benzofuranyl and 2,3-dihydrobenzofuranyl;

where Z^{100} is optionally substituted with one or more substituents each independently selected from the group consisting of F, Cl, CN, optionally substituted alkyl, -O-(optionally substituted alkyl), -COOH, $-Z^{105}-C(O)N(R)_2$, $-Z^{105}-N(R)-C(O)-Z^{200}$, $-Z^{105}-N(R)-S(O)_2-Z^{200}$, and $-Z^{105}-N(R)-C(O)-N(R)-Z^{200}$;

Z^{105} is a covalent bond or (C_1-C_6) ;

Z^{200} is an optionally substituted group selected from group consisting of (C_1-C_6) , phenyl and $-(C_1-C_6)$ -phenyl;


Z^{110} and Z^{111} are each independently a covalent bond or (C_1-C_3) group optionally substituted with alkyl, hydroxy, COOH, CN or phenyl; and

A is O, $-N(R)-C(O)-N(R)-$, $-N(R)-C(O)-O-$, $-N(R)-$ or $-N(R)-C(O)-$, where R is H or alkyl.

-136-


76. A compound according to Claim 75, wherein R_4 is methyl.

77. A compound according to Claim 8, 9 or 10, wherein




78. A compound according to Claim 77, wherein R_4 is methyl; A is -NH-; there is only one R_a and it is H or F; and Z^{100} is optionally substituted with one or more substituents each independently selected from the group consisting of alkyl, halo, CF_3 , and alkoxy.

79. A compound according to Claim 9, wherein



Z^{100} is an optionally substituted group selected from the group consisting of phenyl, pyrrolyl, pyridyl, benzimidazolyl, naphthyl and



where Z^{100} is optionally substituted with one or more substituents each independently selected from the group consisting of F, Cl, Br, NO_2 , amino, N-alkylamino, N,N-dialkylamino, CN, optionally substituted alkyl, -O-(optionally substituted alkyl) and phenyl;


Z^{110} and Z^{111} for each occurrence is independently (C_0-C_3) optionally substituted with optionally substituted phenyl; and

A is -N(R)-C(O)-N(R)-, -N(R)-S(O)₂-, -N(R)-C(O)-, -N(R)- or -N(R)-C(O)-O-.

-137-

80. A compound according to Claim 79, wherein R_4 is methyl and there is only one R_a and it is F.

- 5 81. A compound according to Claim 9 or 66, wherein



Z^{100} is an optionally substituted group selected from the group consisting of phenyl, isoxazolyl, tetrahydronaphthyl, furanyl, benzofuranyl, pyridyl and indolyl;

- 10 where Z^{100} is optionally substituted with one or more substituents each independently selected from the group consisting of F, CN, NO_2 , -C(O)H , -CONH_2 , $\text{-NHSO}_2\text{CF}_3$, optionally substituted alkyl, optionally substituted heteroaryl and $\text{-O-(optionally substituted alkyl)}$;

Z^{110} and Z^{111} are each independently optionally substituted ($\text{C}_0\text{-C}_3$); and


- 15 A is O, $\text{-N(R)-C(O)-(CH}_2\text{)}_n\text{-N(R)-}$, -C(O)-N(R)- , -N(R)-C(O)-O- , -N(R)-C(O)- or -N(R)- .

82. A compound according to Claim 81, wherein R_4 is methyl; R_a is H or methoxy; and Z^{110} and Z^{111} are each unsubstituted.

20


83. A compound according to Claim 9, wherein R_1 is

-138-



where R is H or lower alkyl and n is for each occurrence is independently 1 to 6.


- 5 84. A compound according to Claim 83, wherein R₁ is



85. A compound according to Claim 84, wherein Z¹⁰⁰ is substituted or unsubstituted phenyl.

-139-

86. A compound according to Claim 8, 9 or 10, wherein



5

87. A compound according to Claim 11 wherein n is 2; R_6 is H; m is 1; r is 1; and R_4 and R_5 are each hydrogen.

88. A compound according to claim 64 or 87 wherein R_1 is 4-phenoxyphenyl.

10

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
22 March 2001 (22.03.2001)

PCT

(10) International Publication Number
WO 01/19828 A3

(51) International Patent Classification⁷: **C07D 487/04**,
471/04, 491/04, 498/04, A61K 31/519, 31/5025, 31/4353,
31/551, A61P 35/00 // (C07D 487/04, 239:00, 209:00)
(C07D 487/04, 237:00, 209:00) (C07D 471/04, 231:00,
221:00) (C07D 487/04, 243:00, 239:00)

(GB). **ST. GALLAY, Stephen** [GB/GB]; Bakewell Road,
Loughborough, Leics. LE11 5RH (GB).

(21) International Application Number: **PCT/US00/25357**

(22) International Filing Date:
15 September 2000 (15.09.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/154,618 17 September 1999 (17.09.1999) US

(71) Applicant (for all designated States except US): **BASF AKTIENGESELLSCHAFT** [DE/DE]; 67056 Ludwigshafen (DE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **HIRST, Gavin**,
C. [GB/US]; 112 Robert Road, Marlboro, MA 01752
(US). **RAFFERTY, Paul** [GB/US]; 722 Charlestown
Drive, Westborough, MA 01581 (US). **RITTER, Kurt**
[DE/US]; 30 Ashmont Avenue, Newton, MA 02160 (US).
CALDERWOOD, David [GB/US]; 4 McCarthy Circle,
Framingham, MA 01702 (US). **TWIGGER, Helen, L.**
[GB/GB]; R5 Pennyfoot Street, Nottingham, NG1 1GF

(74) Agents: **WAGNER, Richard, W.** et al.; Hamilton, Brook,
Smith & Reynolds, P.C., Two Militia Drive, Lexington, MA
02421 (US).

(81) Designated States (national): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,
TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG,
CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

(88) Date of publication of the international search report:
4 October 2001

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: **KINASE INHIBITORS AS THERAPEUTIC AGENTS**

(57) Abstract: The present invention is directed to a compound of Formula (I) as defined herein which are useful as kinase inhibitors.

WO 01/19828 A3

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/25357

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D487/04 C07D471/04 C07D491/04 C07D498/04 A61K31/519
A61K31/5025 A61K31/4353 A61K31/551 A61P35/00
//(C07D487/04,239:00,209:00),(C07D487/04,237:00,209:00),

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 98 41525 A (KNOLL AG) 24 September 1998 (1998-09-24) claims 1,28	1,33
A,P	WO 00 44728 A (PFIZER) 3 August 2000 (2000-08-03) claims 1,30	1,33

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- * & * document member of the same patent family

Date of the actual completion of the international search

21 March 2001

Date of mailing of the international search report

29/03/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel (+31-70) 340-2040, Tx. 31 651 epo nl.
Fax: (+31-70) 340-3016

Authorized officer

Alfaro Faus, I

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/25357

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 (C07D471/04, 231:00, 221:00), (C07D487/04, 243:00, 239:00)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- * & * document member of the same patent family

Date of the actual completion of the international search

21 March 2001

Date of mailing of the international search report

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Alfaro Faus, I

INTERNATIONAL SEARCH REPORT

International Application No. PCT/US 00 25357

FURTHER INFORMATION CONTINUED FROM PCT/SA/ 210

Continuation of Box I.2

Present claims 1 to 88 relate to an extremely large number of possible compounds. Support within the meaning of Article 6 PCT and/or disclosure within the meaning of Article 5 PCT is to be found, however, for only a very small proportion of the compounds. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible. Consequently, the search has been carried out for those parts of the claims which appear to be supported and disclosed, namely those parts relating to the compounds of formulas 1 to 117 where R1 is an optionally substituted 4-phenoxyphenyl group and R2 is an optionally substituted cycloalkyl group as described in the examples and on page 46, lines 4 to 6.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 00/25357

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9841525 A	24-09-1998	AU 6829398 A	12-10-1998
		BG 103785 A	30-06-2000
		BR 9808281 A	16-05-2000
		CN 1259950 T	12-07-2000
		EP 0970084 A	12-01-2000
		HU 0001507 A	28-10-2000
		NO 994509 A	17-09-1999
		PL 335685 A	08-05-2000
		SK 125999 A	16-05-2000
		TR 9902301 T	21-12-1999
		US 6001839 A	14-12-1999
WO 0044728 A	03-08-2000	AU 1291600 A	18-08-2000

